

# 1.3

## **SOLVING EQUATIONS WITH VARIABLES ON BOTH SIDES (DAY 1)**

# **DO NOW**

**Solve the equation. Check your solution.**

$$1) \quad \frac{g}{5} - 7 = 12$$

# **DO NOW**

**Solve the equation. Check your solution.**

$$2) \quad 2x + 3x - 5 = 25$$

# **DO NOW**

**Solve the equation. Check your solution.**

$$3) \quad 3(x - 6) + 10 = 16$$

# Solving

- 1) Cancel the “smallest variable term”
- 2) Collect constant terms on the other side

## Examples

$$a) 13 + 5x = 2x - 8$$

$$b) 2m - 6 = 12 - 4m$$

$$c) \quad 34 - 3x = 14x$$

# Practice

1)  $7 - 8x = 4x - 17$

2)  $9 - 3k = 17 - 2k$

## Multi-step with variables on each side of the equation

- 1) Simplify each side of the equation
- 2) Collect variable terms on one side
- 3) Collect constant terms on the other side

### Examples

$$a) 3 - 4y = 5(y - 3)$$

$$b) 3z - 10 + 4z = 5z - 7$$



# Classwork

$$1) y = 24 - 3y$$

# Classwork

$$2) -7a = -12a - 65$$

# Classwork

$$3) 7(a - 2) = 3a + 14$$

# Classwork

$$4) \quad 4(r - 9) + 2 = 12r + 14$$

# Classwork

$$5) 5(2 + n) = 3(n + 6)$$

# Classwork

$$6) 3(2 + v) - 4v = v + 16$$

# No Solution vs Infinitely Many

An equation has **NO SOLUTION**:

if once you solve, one side can NOT be equal to the other side...

An equation is has **INFINITELY MANY SOLUTIONS**:

if once you solve, one side is ALWAYS equal to the other side...

## Examples

a)  $13 + x = 2x - 8$

$$b) \quad 2m - 6 = -6 + 2m$$



$$c) \quad 3x = 3(x + 4)$$