

9 Trigonometric Ratios and Functions

- 9.1 Right Triangle Trigonometry
- 9.2 Angles and Radian Measure
- 9.3 Trigonometric Functions of Any Angle
- 9.4 Graphing Sine and Cosine Functions
- 9.5 Graphing Other Trigonometric Functions
- 9.6 Modeling with Trigonometric Functions
- 9.7 Using Trigonometric Identities
- 9.8 Using Sum and Difference Formulas




Sundial (p. 518)



Tuning Fork (p. 510)



Ferris Wheel (p. 494)



Parasailing (p. 465)



Terminator (p. 476)

Maintaining Mathematical Proficiency

Absolute Value

Example 1 Order the expressions by value from least to greatest: $|6|$, $|-3|$, $\frac{2}{|-4|}$, $|10 - 6|$

$$|6| = 6$$

$$|-3| = 3$$

$$\frac{2}{|-4|} = \frac{2}{4} = \frac{1}{2}$$

$$|10 - 6| = |4| = 4$$

The absolute value of a negative number is positive.

► So, the order is $\frac{2}{|-4|}$, $|-3|$, $|10 - 6|$, and $|6|$.

Order the expressions by value from least to greatest.

1. $|4|$, $|2 - 9|$, $|6 + 4|$, $-|7|$


2. $|9 - 3|$, $|0|$, $|-4|$, $\frac{|-5|}{|2|}$

3. $|-8^3|$, $|-2 \cdot 8|$, $|9 - 1|$, $|9| + |-2| - |1|$

4. $|-4 + 20|$, $-|4^2|$, $|5| - |3 \cdot 2|$, $|-15|$

Pythagorean Theorem

Example 2 Find the missing side length of the triangle.



$$\begin{aligned} a^2 + b^2 &= c^2 \\ 10^2 + b^2 &= 26^2 \\ 100 + b^2 &= 676 \\ b^2 &= 576 \\ b &= 24 \end{aligned}$$

Write the Pythagorean Theorem.

Substitute 10 for a and 26 for c .







Evaluate powers.

Subtract 100 from each side.

Take positive square root of each side.

► So, the length is 24 centimeters.

Find the missing side length of the triangle.



11. **ABSTRACT REASONING** The line segments connecting the points (x_1, y_1) , (x_2, y_1) , and (x_2, y_2) form a triangle. Is the triangle a right triangle? Justify your answer.

Mathematical Practices

Mathematically proficient students reason quantitatively by creating valid representations of problems.

Reasoning Abstractly and Quantitatively


Core Concept

The Unit Circle

The *unit circle* is a circle in the coordinate plane. Its center is at the origin, and it has a radius of 1 unit. The equation of the unit circle is

$$x^2 + y^2 = 1. \quad \text{Equation of unit circle}$$

As the point (x, y) starts at $(1, 0)$ and moves counterclockwise around the unit circle, the angle θ (the Greek letter *theta*) moves from 0° through 360° .



EXAMPLE 1 Finding Coordinates of a Point on the Unit Circle

Find the exact coordinates of the point (x, y) on the unit circle.

SOLUTION

Because $\theta = 45^\circ$, (x, y) lies on the line $y = x$.

$$x^2 + y^2 = 1$$

Write equation of unit circle.

$$x^2 + x^2 = 1$$

Substitute x for y .

$$2x^2 = 1$$


Add like terms.

$$x^2 = \frac{1}{2}$$

Divide each side by 2.

$$x = \frac{1}{\sqrt{2}}$$

Take positive square root of each side.




► The coordinates of (x, y) are $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$, or $\left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$.


Monitoring Progress

Find the exact coordinates of the point (x, y) on the unit circle.


1.



2.



3.




9.1 Right Triangle Trigonometry

Essential Question How can you find a trigonometric function of an acute angle θ ?



Consider one of the acute angles θ of a right triangle. Ratios of a right triangle's side lengths are used to define the six *trigonometric functions*, as shown.

Sine	$\sin \theta = \frac{\text{opp.}}{\text{hyp.}}$	Cosine	$\cos \theta = \frac{\text{adj.}}{\text{hyp.}}$
Tangent	$\tan \theta = \frac{\text{opp.}}{\text{adj.}}$	Cotangent	$\cot \theta = \frac{\text{adj.}}{\text{opp.}}$
Secant	$\sec \theta = \frac{\text{hyp.}}{\text{adj.}}$	Cosecant	$\csc \theta = \frac{\text{hyp.}}{\text{opp.}}$



EXPLORATION 1 Trigonometric Functions of Special Angles

Work with a partner. Find the exact values of the sine, cosine, and tangent functions for the angles 30° , 45° , and 60° in the right triangles shown.



CONSTRUCTING VIABLE ARGUMENTS

To be proficient in math, you need to understand and use stated assumptions, definitions, and previously established results in constructing arguments.

EXPLORATION 2 Exploring Trigonometric Identities

Work with a partner.

Use the definitions of the trigonometric functions to explain why each *trigonometric identity* is true.

a. $\sin \theta = \cos(90^\circ - \theta)$

b. $\cos \theta = \sin(90^\circ - \theta)$

c. $\sin \theta = \frac{1}{\csc \theta}$

d. $\tan \theta = \frac{1}{\cot \theta}$

Use the definitions of the trigonometric functions to complete each trigonometric identity.


e. $(\sin \theta)^2 + (\cos \theta)^2 =$

f. $(\sec \theta)^2 - (\tan \theta)^2 =$

Communicate Your Answer

3. How can you find a trigonometric function of an acute angle θ ?

4. Use a calculator to find the lengths x and y of the legs of the right triangle shown.



9.1 Lesson

Core Vocabulary

sine, p. 462
cosine, p. 462
tangent, p. 462
cosecant, p. 462
secant, p. 462
cotangent, p. 462

Previous

right triangle
hypotenuse
acute angle
Pythagorean Theorem
reciprocal
complementary angles


What You Will Learn

- ▶ Evaluate trigonometric functions of acute angles.
- ▶ Find unknown side lengths and angle measures of right triangles.
- ▶ Use trigonometric functions to solve real-life problems.

The Six Trigonometric Functions

Consider a right triangle that has an acute angle θ (the Greek letter *theta*). The three sides of the triangle are the *hypotenuse*, the side *opposite* θ , and the side *adjacent* to θ .

Ratios of a right triangle's side lengths are used to define the six trigonometric functions: **sine**, **cosine**, **tangent**, **cosecant**, **secant**, and **cotangent**. These six functions are abbreviated \sin , \cos , \tan , \csc , \sec , and \cot , respectively.



Core Concept

Right Triangle Definitions of Trigonometric Functions

Let θ be an acute angle of a right triangle. The six trigonometric functions of θ are defined as shown.


$$\begin{aligned}\sin \theta &= \frac{\text{opposite}}{\text{hypotenuse}} & \cos \theta &= \frac{\text{adjacent}}{\text{hypotenuse}} & \tan \theta &= \frac{\text{opposite}}{\text{adjacent}} \\ \csc \theta &= \frac{\text{hypotenuse}}{\text{opposite}} & \sec \theta &= \frac{\text{hypotenuse}}{\text{adjacent}} & \cot \theta &= \frac{\text{adjacent}}{\text{opposite}}\end{aligned}$$

The abbreviations *opp.*, *adj.*, and *hyp.* are often used to represent the side lengths of the right triangle. Note that the ratios in the second row are reciprocals of the ratios in the first row.

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

REMEMBER

The Pythagorean Theorem states that $a^2 + b^2 = c^2$ for a right triangle with hypotenuse of length c and legs of lengths a and b .



EXAMPLE 1 Evaluating Trigonometric Functions

Evaluate the six trigonometric functions of the angle θ .


SOLUTION

From the Pythagorean Theorem, the length of the hypotenuse is

$$\begin{aligned}\text{hyp.} &= \sqrt{5^2 + 12^2} \\ &= \sqrt{169} \\ &= 13.\end{aligned}$$

Using $\text{adj.} = 5$, $\text{opp.} = 12$, and $\text{hyp.} = 13$, the values of the six trigonometric functions of θ are:

$$\begin{aligned}\sin \theta &= \frac{\text{opp.}}{\text{hyp.}} = \frac{12}{13} & \cos \theta &= \frac{\text{adj.}}{\text{hyp.}} = \frac{5}{13} & \tan \theta &= \frac{\text{opp.}}{\text{adj.}} = \frac{12}{5} \\ \csc \theta &= \frac{\text{hyp.}}{\text{opp.}} = \frac{13}{12} & \sec \theta &= \frac{\text{hyp.}}{\text{adj.}} = \frac{13}{5} & \cot \theta &= \frac{\text{adj.}}{\text{opp.}} = \frac{5}{12}\end{aligned}$$




EXAMPLE 2 Evaluating Trigonometric Functions

In a right triangle, θ is an acute angle and $\sin \theta = \frac{4}{7}$. Evaluate the other five trigonometric functions of θ .

SOLUTION

Step 1 Draw a right triangle with acute angle θ such that the leg opposite θ has length 4 and the hypotenuse has length 7.



Step 2 Find the length of the adjacent side. By the Pythagorean Theorem, the length of the other leg is

$$\text{adj.} = \sqrt{7^2 - 4^2} = \sqrt{33}.$$

Step 3 Find the values of the remaining five trigonometric functions.

Because $\sin \theta = \frac{4}{7}$, $\csc \theta = \frac{\text{hyp.}}{\text{opp.}} = \frac{7}{4}$. The other values are:

$$\cos \theta = \frac{\text{adj.}}{\text{hyp.}} = \frac{\sqrt{33}}{7} \qquad \tan \theta = \frac{\text{opp.}}{\text{adj.}} = \frac{4}{\sqrt{33}} = \frac{4\sqrt{33}}{33}$$

$$\sec \theta = \frac{\text{hyp.}}{\text{adj.}} = \frac{7}{\sqrt{33}} = \frac{7\sqrt{33}}{33} \qquad \cot \theta = \frac{\text{adj.}}{\text{opp.}} = \frac{\sqrt{33}}{4}$$


Monitoring Progress




Help in English and Spanish at BigIdeasMath.com

Evaluate the six trigonometric functions of the angle θ .


1.



2.



3.




4. In a right triangle, θ is an acute angle and $\cos \theta = \frac{7}{10}$. Evaluate the other five trigonometric functions of θ .

The angles 30° , 45° , and 60° occur frequently in trigonometry. You can use the trigonometric values for these angles to find unknown side lengths in special right triangles.

Core Concept

Trigonometric Values for Special Angles

The table gives the values of the six trigonometric functions for the angles 30° , 45° , and 60° . You can obtain these values from the triangles shown.




θ	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\csc \theta$	$\sec \theta$	$\cot \theta$
30°	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$	$\sqrt{3}$
45°	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	$\sqrt{2}$	$\sqrt{2}$	1
60°	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2	$\frac{\sqrt{3}}{3}$

Finding Side Lengths and Angle Measures

EXAMPLE 3 Finding an Unknown Side Length

Find the value of x for the right triangle.



SOLUTION

Write an equation using a trigonometric function that involves the ratio of x and 8. Solve the equation for x .

$$\cos 30^\circ = \frac{\text{adj.}}{\text{hyp.}} \quad \text{Write trigonometric equation.}$$

$$\frac{\sqrt{3}}{2} = \frac{x}{8} \quad \text{Substitute.}$$

$$4\sqrt{3} = x \quad \text{Multiply each side by 8.}$$

► The length of the side is $x = 4\sqrt{3} \approx 6.93$.

READING

Throughout this chapter, a capital letter is used to denote both an angle of a triangle and its measure. The same letter in lowercase is used to denote the length of the side opposite that angle.


Finding all unknown side lengths and angle measures of a triangle is called *solving the triangle*. Solving right triangles that have acute angles other than 30° , 45° , and 60° may require the use of a calculator. Be sure the calculator is set in *degree* mode.

EXAMPLE 4 Using a Calculator to Solve a Right Triangle

Solve $\triangle ABC$.

SOLUTION

Because the triangle is a right triangle, A and B are complementary angles. So, $B = 90^\circ - 28^\circ = 62^\circ$.



Next, write two equations using trigonometric functions, one that involves the ratio of a and 15, and one that involves c and 15. Solve the first equation for a and the second equation for c .

$$\tan 28^\circ = \frac{\text{opp.}}{\text{adj.}} \quad \text{Write trigonometric equation.} \quad \sec 28^\circ = \frac{\text{hyp.}}{\text{adj.}}$$

$$\tan 28^\circ = \frac{a}{15} \quad \text{Substitute.} \quad \sec 28^\circ = \frac{c}{15}$$



$$15(\tan 28^\circ) = a \quad \text{Solve for the variable.} \quad 15\left(\frac{1}{\cos 28^\circ}\right) = c$$

$$7.98 \approx a \quad \text{Use a calculator.} \quad 16.99 \approx c$$

► So, $B = 62^\circ$, $a \approx 7.98$, and $c \approx 16.99$.

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

5. Find the value of x for the right triangle shown.



Solve $\triangle ABC$ using the diagram at the left and the given measurements.

6. $B = 45^\circ$, $c = 5$

7. $A = 32^\circ$, $b = 10$

8. $A = 71^\circ$, $c = 20$

9. $B = 60^\circ$, $a = 7$

Solving Real-Life Problems

EXAMPLE 5 Using Indirect Measurement

FINDING AN ENTRY POINT

The tangent function is used to find the unknown distance because it involves the ratio of x and 2.

You are hiking near a canyon. While standing at A , you measure an angle of 90° between B and C , as shown. You then walk to B and measure an angle of 76° between A and C . The distance between A and B is about 2 miles. How wide is the canyon between A and C ?

SOLUTION

$$\tan 76^\circ = \frac{x}{2}$$

Write trigonometric equation.


$$2(\tan 76^\circ) = x$$

Multiply each side by 2.


$$8.0 \approx x$$

Use a calculator.

► The width is about 8.0 miles.



If you look at a point above you, such as the top of a building, the angle that your line of sight makes with a line parallel to the ground is called the *angle of elevation*. At the top of the building, the angle between a line parallel to the ground and your line of sight is called the *angle of depression*. These two angles have the same measure.




EXAMPLE 6 Using an Angle of Elevation

A parasailer is attached to a boat with a rope 72 feet long. The angle of elevation from the boat to the parasailer is 28° . Estimate the parasailer's height above the boat.

SOLUTION

Step 1 Draw a diagram that represents the situation.



Step 2 Write and solve an equation to find the height h .

$$\sin 28^\circ = \frac{h}{72}$$

Write trigonometric equation.

$$72(\sin 28^\circ) = h$$

Multiply each side by 72.

$$33.8 \approx h$$

Use a calculator.

► The height of the parasailer above the boat is about 33.8 feet.



Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

10. In Example 5, find the distance between B and C .

11. **WHAT IF?** In Example 6, estimate the height of the parasailer above the boat when the angle of elevation is 38° .

9.1 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check


- COMPLETE THE SENTENCE** In a right triangle, the two trigonometric functions of θ that are defined using the lengths of the hypotenuse and the side adjacent to θ are _____ and _____.
- VOCABULARY** Compare an angle of elevation to an angle of depression.
- WRITING** Explain what it means to solve a right triangle.
- DIFFERENT WORDS, SAME QUESTION** Which is different? Find “both” answers.

What is the cosecant of θ ?

What is $\frac{1}{\sin \theta}$?







What is the ratio of the side opposite θ to the hypotenuse?

What is the ratio of the hypotenuse to the side opposite θ ?



Monitoring Progress and Modeling with Mathematics

In Exercises 5–10, evaluate the six trigonometric functions of the angle θ . (See Example 1.)



11. **REASONING** Let θ be an acute angle of a right triangle. Use the two trigonometric functions $\tan \theta = \frac{4}{9}$ and $\sec \theta = \frac{\sqrt{97}}{9}$ to sketch and label the right triangle. Then evaluate the other four trigonometric functions of θ .

12. **ANALYZING RELATIONSHIPS** Evaluate the six trigonometric functions of the $90^\circ - \theta$ angle in Exercises 5–10. Describe the relationships you notice.

In Exercises 13–18, let θ be an acute angle of a right triangle. Evaluate the other five trigonometric functions of θ . (See Example 2.)

13. $\sin \theta = \frac{7}{11}$

14. $\cos \theta = \frac{5}{12}$


15. $\tan \theta = \frac{7}{6}$

16. $\csc \theta = \frac{15}{8}$

17. $\sec \theta = \frac{14}{9}$

18. $\cot \theta = \frac{16}{11}$

19. **ERROR ANALYSIS** Describe and correct the error in finding $\sin \theta$ of the triangle below.









$$\sin \theta = \frac{\text{opp.}}{\text{hyp.}} = \frac{15}{17}$$

20. **ERROR ANALYSIS** Describe and correct the error in finding $\csc \theta$, given that θ is an acute angle of a right triangle and $\cos \theta = \frac{7}{11}$.



$$\csc \theta = \frac{1}{\cos \theta} = \frac{11}{7}$$


In Exercises 21–26, find the value of x for the right triangle. (See Example 3.)



USING TOOLS In Exercises 27–32, evaluate the trigonometric function using a calculator. Round your answer to four decimal places.


27. $\cos 14^\circ$ 28. $\tan 31^\circ$
 29. $\csc 59^\circ$ 30. $\sin 23^\circ$
 31. $\cot 6^\circ$ 32. $\sec 11^\circ$

In Exercises 33–40, solve $\triangle ABC$ using the diagram and the given measurements. (See Example 4.)




33. $B = 36^\circ$, $a = 23$ 34. $A = 27^\circ$, $b = 9$
 35. $A = 55^\circ$, $a = 17$ 36. $B = 16^\circ$, $b = 14$
 37. $A = 43^\circ$, $b = 31$ 38. $B = 31^\circ$, $a = 23$
 39. $B = 72^\circ$, $c = 12.8$ 40. $A = 64^\circ$, $a = 7.4$

41. **MODELING WITH MATHEMATICS** To measure the width of a river, you plant a stake on one side of the river, directly across from a boulder. You then walk 100 meters to the right of the stake and measure a 79° angle between the stake and the boulder. What is the width w of the river? (See Example 5.)




42. **MODELING WITH MATHEMATICS** Katoomba Scenic Railway in Australia is the steepest railway in the world. The railway makes an angle of about 52° with the ground. The railway extends horizontally about 458 feet. What is the height of the railway?
43. **MODELING WITH MATHEMATICS** A person whose eye level is 1.5 meters above the ground is standing 75 meters from the base of the Jin Mao Building in Shanghai, China. The person estimates the angle of elevation to the top of the building is about 80° . What is the approximate height of the building? (See Example 6.)
44. **MODELING WITH MATHEMATICS** The Duquesne Incline in Pittsburgh, Pennsylvania, has an angle of elevation of 30° . The track has a length of about 800 feet. Find the height of the incline.
45. **MODELING WITH MATHEMATICS** You are standing on the Grand View Terrace viewing platform at Mount Rushmore, 1000 feet from the base of the monument.




- a. You look up at the top of Mount Rushmore at an angle of 24° . How high is the top of the monument from where you are standing? Assume your eye level is 5.5 feet above the platform.
- b. The elevation of the Grand View Terrace is 5280 feet. Use your answer in part (a) to find the elevation of the top of Mount Rushmore.
46. **WRITING** Write a real-life problem that can be solved using a right triangle. Then solve your problem.

- 47. MATHEMATICAL CONNECTIONS** The Tropic of Cancer is the circle of latitude farthest north of the equator where the Sun can appear directly overhead. It lies 23.5° north of the equator, as shown.




- Find the circumference of the Tropic of Cancer using 3960 miles as the approximate radius of Earth.
- What is the distance between two points on the Tropic of Cancer that lie directly across from each other?

- 48. HOW DO YOU SEE IT?** Use the figure to answer each question.



- Which side is adjacent to θ ?
- Which side is opposite of θ ?
- Does $\cos \theta = \sin(90^\circ - \theta)$? Explain.

- 49. PROBLEM SOLVING** A passenger in an airplane sees two towns directly to the left of the plane.




- What is the distance d from the airplane to the first town?
- What is the horizontal distance x from the airplane to the first town?
- What is the distance y between the two towns? Explain the process you used to find your answer.


- 50. PROBLEM SOLVING** You measure the angle of elevation from the ground to the top of a building as 32° . When you move 50 meters closer to the building, the angle of elevation is 53° . What is the height of the building?

- 51. MAKING AN ARGUMENT** Your friend claims it is possible to draw a right triangle so the values of the cosine function of the acute angles are equal. Is your friend correct? Explain your reasoning.

- 52. THOUGHT PROVOKING** Consider a semicircle with a radius of 1 unit, as shown below. Write the values of the six trigonometric functions of the angle θ . Explain your reasoning.



- 53. CRITICAL THINKING** A procedure for approximating π based on the work of Archimedes is to inscribe a regular hexagon in a circle.



- Use the diagram to solve for x . What is the perimeter of the hexagon?
- Show that a regular n -sided polygon inscribed in a circle of radius 1 has a perimeter of $2n \cdot \sin\left(\frac{180}{n}\right)^\circ$.
- Use the result from part (b) to find an expression in terms of n that approximates π . Then evaluate the expression when $n = 50$.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Perform the indicated conversion. (*Skills Review Handbook*)

54. 5 years to seconds

55. 12 pints to gallons

56. 5.6 meters to millimeters

Find the circumference and area of the circle with the given radius or diameter.

(*Skills Review Handbook*)

57. $r = 6$ centimeters

58. $r = 11$ inches

59. $d = 14$ feet

9.2 Angles and Radian Measure


Essential Question How can you find the measure of an angle in radians?

Let the vertex of an angle be at the origin, with one side of the angle on the positive x -axis. The *radian measure* of the angle is a measure of the intercepted arc length on a circle of radius 1. To convert between degree and radian measure, use the fact that

$$\frac{\pi \text{ radians}}{180^\circ} = 1.$$


EXPLORATION 1 Writing Radian Measures of Angles

Work with a partner. Write the radian measure of each angle with the given degree measure. Explain your reasoning.



EXPLORATION 2 Writing Degree Measures of Angles

Work with a partner. Write the degree measure of each angle with the given radian measure. Explain your reasoning.




REASONING ABSTRACTLY

To be proficient in math, you need to make sense of quantities and their relationships in problem situations.

Communicate Your Answer

- How can you find the measure of an angle in radians?
- The figure shows an angle whose measure is 30 radians. What is the measure of the angle in degrees? How many times greater is 30 radians than 30 degrees? Justify your answers.



9.2 Lesson

Core Vocabulary

initial side, p. 470
terminal side, p. 470
standard position, p. 470
coterminal, p. 471
radian, p. 471
sector, p. 472
central angle, p. 472

Previous

radius of a circle
circumference of a circle

What You Will Learn

- ▶ Draw angles in standard position.
- ▶ Find coterminal angles.
- ▶ Use radian measure.

Drawing Angles in Standard Position


In this lesson, you will expand your study of angles to include angles with measures that can be any real numbers.

Core Concept

Angles in Standard Position

In a coordinate plane, an angle can be formed by fixing one ray, called the **initial side**, and rotating the other ray, called the **terminal side**, about the vertex.

An angle is in **standard position** when its vertex is at the origin and its initial side lies on the positive x -axis.



The measure of an angle is positive when the rotation of its terminal side is counterclockwise and negative when the rotation is clockwise. The terminal side of an angle can rotate more than 360° .

EXAMPLE 1

Drawing Angles in Standard Position

Draw an angle with the given measure in standard position.


a. 240°

b. 500°


c. -50°

SOLUTION


a. Because 240° is 60° more than 180° , the terminal side is 60° counterclockwise past the negative x -axis.



b. Because 500° is 140° more than 360° , the terminal side makes one complete rotation 360° counterclockwise plus 140° more.



c. Because -50° is negative, the terminal side is 50° clockwise from the positive x -axis.



Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Draw an angle with the given measure in standard position.

1. 65°

2. 300°

3. -120°

4. -450°

STUDY TIP

If two angles differ by a multiple of 360° , then the angles are coterminal.

Finding Coterminal Angles

In Example 1(b), the angles 500° and 140° are **coterminal** because their terminal sides coincide. An angle coterminal with a given angle can be found by adding or subtracting multiples of 360° .

EXAMPLE 2

Finding Coterminal Angles



Find one positive angle and one negative angle that are coterminal with (a) -45° and (b) 395° .

SOLUTION

There are many such angles, depending on what multiple of 360° is added or subtracted.

a. $-45^\circ + 360^\circ = 315^\circ$
 $-45^\circ - 360^\circ = -405^\circ$

b. $395^\circ - 360^\circ = 35^\circ$
 $395^\circ - 2(360^\circ) = -325^\circ$



Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Find one positive angle and one negative angle that are coterminal with the given angle.

5. 80°

6. 230°

7. 740°

8. -135°

STUDY TIP

Notice that 1 radian is approximately equal to 57.3° .

$$180^\circ = \pi \text{ radians}$$


$$\frac{180^\circ}{\pi} = 1 \text{ radian}$$

$$57.3^\circ \approx 1 \text{ radian}$$

Using Radian Measure

Angles can also be measured in *radians*. To define a radian, consider a circle with radius r centered at the origin, as shown. One **radian** is the measure of an angle in standard position whose terminal side intercepts an arc of length r .

Because the circumference of a circle is $2\pi r$, there are 2π radians in a full circle. So, degree measure and radian measure are related by the equation $360^\circ = 2\pi$ radians, or $180^\circ = \pi$ radians.



Core Concept

Converting Between Degrees and Radians

Degrees to radians

Multiply degree measure by

$$\frac{\pi \text{ radians}}{180^\circ}$$

Radians to degrees

Multiply radian measure by

$$\frac{180^\circ}{\pi \text{ radians}}$$

EXAMPLE 3**Convert Between Degrees and Radians**

Convert the degree measure to radians or the radian measure to degrees.

READING

The unit “radians” is often omitted. For instance, the measure $-\frac{\pi}{12}$ radians may be written simply as $-\frac{\pi}{12}$.

a. 120°

b. $-\frac{\pi}{12}$

SOLUTION


$$\begin{aligned} \text{a. } 120^\circ &= 120 \text{ degrees} \left(\frac{\pi \text{ radians}}{180 \text{ degrees}} \right) \\ &= \frac{2\pi}{3} \end{aligned}$$

$$\begin{aligned} \text{b. } -\frac{\pi}{12} &= \left(-\frac{\pi}{12} \text{ radians} \right) \left(\frac{180^\circ}{\pi \text{ radians}} \right) \\ &= -15^\circ \end{aligned}$$

Concept Summary**Degree and Radian Measures of Special Angles**

The diagram shows equivalent degree and radian measures for special angles from 0° to 360° (0 radians to 2π radians).

You may find it helpful to memorize the equivalent degree and radian measures of special angles in the first quadrant and for $90^\circ = \frac{\pi}{2}$ radians. All other special angles shown are multiples of these angles.

**Monitoring Progress**

Help in English and Spanish at BigIdeasMath.com

Convert the degree measure to radians or the radian measure to degrees.

9. 135°

10. -40°

11. $\frac{5\pi}{4}$

12. -6.28


A **sector** is a region of a circle that is bounded by two radii and an arc of the circle. The **central angle** θ of a sector is the angle formed by the two radii. There are simple formulas for the arc length and area of a sector when the central angle is measured in radians.

Core Concept**Arc Length and Area of a Sector**

The arc length s and area A of a sector with radius r and central angle θ (measured in radians) are as follows.

Arc length: $s = r\theta$

Area: $A = \frac{1}{2}r^2\theta$




EXAMPLE 4

Modeling with Mathematics

A softball field forms a sector with the dimensions shown. Find the length of the outfield fence and the area of the field.

SOLUTION

- 1. Understand the Problem** You are given the dimensions of a softball field. You are asked to find the length of the outfield fence and the area of the field.
- 2. Make a Plan** Find the measure of the central angle in radians. Then use the arc length and area of a sector formulas.
- 3. Solve the Problem**



Step 1 Convert the measure of the central angle to radians.

$$\begin{aligned} 90^\circ &= 90 \text{ degrees} \left(\frac{\pi \text{ radians}}{180 \text{ degrees}} \right) \\ &= \frac{\pi}{2} \text{ radians} \end{aligned}$$

Step 2 Find the arc length and the area of the sector.

Arc length: $s = r\theta$

$$\begin{aligned} &= 200 \left(\frac{\pi}{2} \right) \\ &= 100\pi \\ &\approx 314 \end{aligned}$$


Area: $A = \frac{1}{2}r^2\theta$

$$\begin{aligned} &= \frac{1}{2}(200)^2 \left(\frac{\pi}{2} \right) \\ &= 10,000\pi \\ &\approx 31,416 \end{aligned}$$

► The length of the outfield fence is about 314 feet. The area of the field is about 31,416 square feet.

- 4. Look Back** To check the area of the field, consider the square formed using the two 200-foot sides.

By drawing the diagonal, you can see that the area of the field is less than the area of the square but greater than one-half of the area of the square.



$\frac{1}{2} \cdot (\text{area of square})$

area of square

$$\frac{1}{2}(200)^2 \stackrel{?}{<} 31,416 \stackrel{?}{<} 200^2$$

$$20,000 < 31,416 < 40,000 \quad \checkmark$$

COMMON ERROR

You must write the measure of an angle in radians when using these formulas for the arc length and area of a sector.

ANOTHER WAY

Because the central angle is 90° , the sector represents $\frac{1}{4}$ of a circle with a radius of 200 feet. So,

$$\begin{aligned} s &= \frac{1}{4} \cdot 2\pi r = \frac{1}{4} \cdot 2\pi(200) \\ &= 100\pi \end{aligned}$$

and

$$\begin{aligned} A &= \frac{1}{4} \cdot \pi r^2 = \frac{1}{4} \cdot \pi(200)^2 \\ &= 10,000\pi. \end{aligned}$$

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

- 13. WHAT IF?** In Example 4, the outfield fence is 220 feet from home plate. Estimate the length of the outfield fence and the area of the field.

9.2 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** An angle is in standard position when its vertex is at the _____ and its _____ lies on the positive x -axis.
- WRITING** Explain how the sign of an angle measure determines its direction of rotation.
- VOCABULARY** In your own words, define a radian.
- WHICH ONE DOESN'T BELONG?** Which angle does *not* belong with the other three? Explain your reasoning.

-90°

450°

90°

-270°

Monitoring Progress and Modeling with Mathematics

In Exercises 5–8, draw an angle with the given measure in standard position. (See Example 1.)

- 110°
- 450°
- -900°
- -10°

In Exercises 9–12, find one positive angle and one negative angle that are coterminal with the given angle. (See Example 2.)


- 70°
- 255°
- -125°
- -800°

In Exercises 13–20, convert the degree measure to radians or the radian measure to degrees. (See Example 3.)

- 40°
- 315°
- -260°
- -500°
- $\frac{\pi}{9}$
- $\frac{3\pi}{4}$
- -5
- 12

- WRITING** The terminal side of an angle in standard position rotates one-sixth of a revolution counterclockwise from the positive x -axis. Describe how to find the measure of the angle in both degree and radian measures.


- OPEN-ENDED** Using radian measure, give one positive angle and one negative angle that are coterminal with the angle shown. Justify your answers.




ANALYZING RELATIONSHIPS In Exercises 23–26, match the angle measure with the angle.

- 600°
- $-\frac{9\pi}{4}$
- $\frac{5\pi}{6}$
- -240°


A.




B.




C.



D.



27. **MODELING WITH MATHEMATICS** The observation deck of a building forms a sector with the dimensions shown. Find the length of the safety rail and the area of the deck. (See Example 4.)



28. **MODELING WITH MATHEMATICS** In the men's shot put event at the 2012 Summer Olympic Games, the length of the winning shot was 21.89 meters. A shot put must land within a sector having a central angle of 34.92° to be considered fair.



- The officials draw an arc across the fair landing area, marking the farthest throw. Find the length of the arc.
 - All fair throws in the 2012 Olympics landed within a sector bounded by the arc in part (a). What is the area of this sector?
29. **ERROR ANALYSIS** Describe and correct the error in converting the degree measure to radians.

$$\begin{aligned}
 24^\circ &= 24 \text{ ~~degrees~~ } \left(\frac{180 \text{ ~~degrees~~}}{\pi \text{ radians}} \right) \\
 &= \frac{4320}{\pi} \text{ radians} \\
 &\approx 1375.1 \text{ radians}
 \end{aligned}$$

30. **ERROR ANALYSIS** Describe and correct the error in finding the area of a sector with a radius of 6 centimeters and a central angle of 40° .

$$A = \frac{1}{2}(6)^2(40) = 720 \text{ cm}^2$$

31. **PROBLEM SOLVING** When a CD player reads information from the outer edge of a CD, the CD spins about 200 revolutions per minute. At that speed, through what angle does a point on the CD spin in one minute? Give your answer in both degree and radian measures.

32. **PROBLEM SOLVING** You work every Saturday from 9:00 A.M. to 5:00 P.M. Draw a diagram that shows the rotation completed by the hour hand of a clock during this time. Find the measure of the angle generated by the hour hand in both degrees and radians. Compare this angle with the angle generated by the minute hand from 9:00 A.M. to 5:00 P.M.

USING TOOLS In Exercises 33–38, use a calculator to evaluate the trigonometric function.

33. $\cos \frac{4\pi}{3}$

34. $\sin \frac{7\pi}{8}$


35. $\csc \frac{10\pi}{11}$

36. $\cot\left(-\frac{6\pi}{5}\right)$

37. $\cot(-14)$

38. $\cos 6$

39. **MODELING WITH MATHEMATICS** The rear windshield wiper of a car rotates 120° , as shown. Find the area cleared by the wiper.



40. **MODELING WITH MATHEMATICS** A scientist performed an experiment to study the effects of gravitational force on humans. In order for humans to experience twice Earth's gravity, they were placed in a centrifuge 58 feet long and spun at a rate of about 15 revolutions per minute.




- Through how many radians did the people rotate each second?
- Find the length of the arc through which the people rotated each second.


41. **REASONING** In astronomy, the *terminator* is the day-night line on a planet that divides the planet into daytime and nighttime regions. The terminator moves across the surface of a planet as the planet rotates. It takes about 4 hours for Earth's terminator to move across the continental United States. Through what angle has Earth rotated during this time? Give your answer in both degree and radian measures.



42. **HOW DO YOU SEE IT?** Use the graph to find the measure of θ . Explain your reasoning.



43. **MODELING WITH MATHEMATICS** A dartboard is divided into 20 sectors. Each sector is worth a point value from 1 to 20 and has shaded regions that double or triple this value. A sector is shown below. Find the areas of the entire sector, the double region, and the triple region.



44. **THOUGHT PROVOKING** π is an irrational number, which means that it cannot be written as the ratio of two whole numbers. π can, however, be written exactly as a *continued fraction*, as follows.

$$3 + \frac{1}{7 + \frac{1}{15 + \frac{1}{1 + \frac{1}{292 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}}}}}}$$

Show how to use this continued fraction to obtain a decimal approximation for π .

45. **MAKING AN ARGUMENT** Your friend claims that when the arc length of a sector equals the radius, the area can be given by $A = \frac{s^2}{2}$. Is your friend correct? Explain.
46. **PROBLEM SOLVING** A spiral staircase has 15 steps. Each step is a sector with a radius of 42 inches and a central angle of $\frac{\pi}{8}$.
- What is the length of the arc formed by the outer edge of a step?
 - Through what angle would you rotate by climbing the stairs?
 - How many square inches of carpeting would you need to cover the 15 steps?
47. **MULTIPLE REPRESENTATIONS** There are 60 *minutes* in 1 degree of arc, and 60 *seconds* in 1 minute of arc. The notation $50^\circ 30' 10''$ represents an angle with a measure of 50 degrees, 30 minutes, and 10 seconds.
- Write the angle measure 70.55° using the notation above.
 - Write the angle measure $110^\circ 45' 30''$ to the nearest hundredth of a degree. Justify your answer.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Find the distance between the two points. (*Skills Review Handbook*)

- | | |
|----------------------------|------------------------|
| 48. (1, 4), (3, 6) | 49. (-7, -13), (10, 8) |
| 50. (-3, 9), (-3, 16) | 51. (2, 12), (8, -5) |
| 52. (-14, -22), (-20, -32) | 53. (4, 16), (-1, 34) |

9.3 Trigonometric Functions of Any Angle


Essential Question How can you use the unit circle to define the trigonometric functions of any angle?

Let θ be an angle in standard position with (x, y) a point on the terminal side of θ and $r = \sqrt{x^2 + y^2} \neq 0$. The six trigonometric functions of θ are defined as shown.

$$\sin \theta = \frac{y}{r} \qquad \csc \theta = \frac{r}{y}, y \neq 0$$







$$\cos \theta = \frac{x}{r} \qquad \sec \theta = \frac{r}{x}, x \neq 0$$

$$\tan \theta = \frac{y}{x}, x \neq 0 \qquad \cot \theta = \frac{x}{y}, y \neq 0$$



EXPLORATION 1 Writing Trigonometric Functions

Work with a partner. Find the sine, cosine, and tangent of the angle θ in standard position whose terminal side intersects the unit circle at the point (x, y) shown.



CONSTRUCTING VIABLE ARGUMENTS

To be proficient in math, you need to understand and use stated assumptions, definitions, and previously established results.

Communicate Your Answer

- How can you use the unit circle to define the trigonometric functions of any angle?
- For which angles are each function undefined? Explain your reasoning.
 - tangent
 - cotangent
 - secant
 - cosecant

9.3 Lesson

Core Vocabulary

unit circle, p. 479
 quadrantal angle, p. 479
 reference angle, p. 480

Previous

circle
 radius
 Pythagorean Theorem

What You Will Learn

- ▶ Evaluate trigonometric functions of any angle.
- ▶ Find and use reference angles to evaluate trigonometric functions.

Trigonometric Functions of Any Angle

You can generalize the right-triangle definitions of trigonometric functions so that they apply to any angle in standard position.

Core Concept

General Definitions of Trigonometric Functions

Let θ be an angle in standard position, and let (x, y) be the point where the terminal side of θ intersects the circle $x^2 + y^2 = r^2$. The six trigonometric functions of θ are defined as shown.

$$\sin \theta = \frac{y}{r}$$

$$\csc \theta = \frac{r}{y}, y \neq 0$$


$$\cos \theta = \frac{x}{r}$$

$$\sec \theta = \frac{r}{x}, x \neq 0$$

$$\tan \theta = \frac{y}{x}, x \neq 0$$

$$\cot \theta = \frac{x}{y}, y \neq 0$$

These functions are sometimes called *circular functions*.



EXAMPLE 1

Evaluating Trigonometric Functions Given a Point

Let $(-4, 3)$ be a point on the terminal side of an angle θ in standard position. Evaluate the six trigonometric functions of θ .

SOLUTION

Use the Pythagorean Theorem to find the length of r .

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{(-4)^2 + 3^2} \\ &= \sqrt{25} \\ &= 5 \end{aligned}$$

Using $x = -4$, $y = 3$, and $r = 5$, the values of the six trigonometric functions of θ are:

$$\sin \theta = \frac{y}{r} = \frac{3}{5}$$


$$\csc \theta = \frac{r}{y} = \frac{5}{3}$$

$$\cos \theta = \frac{x}{r} = -\frac{4}{5}$$

$$\sec \theta = \frac{r}{x} = -\frac{5}{4}$$

$$\tan \theta = \frac{y}{x} = -\frac{3}{4}$$

$$\cot \theta = \frac{x}{y} = -\frac{4}{3}$$




Core Concept

The Unit Circle

The circle $x^2 + y^2 = 1$, which has center $(0, 0)$ and radius 1, is called the **unit circle**. The values of $\sin \theta$ and $\cos \theta$ are simply the y -coordinate and x -coordinate, respectively, of the point where the terminal side of θ intersects the unit circle.

$$\sin \theta = \frac{y}{r} = \frac{y}{1} = y$$

$$\cos \theta = \frac{x}{r} = \frac{x}{1} = x$$



ANOTHER WAY

The general circle $x^2 + y^2 = r^2$ can also be used to find the six trigonometric functions of θ . The terminal side of θ intersects the circle at $(0, -r)$. So,

$$\sin \theta = \frac{y}{r} = \frac{-r}{r} = -1.$$

The other functions can be evaluated similarly.

It is convenient to use the unit circle to find trigonometric functions of **quadrantal angles**. A quadrantal angle is an angle in standard position whose terminal side lies on an axis. The measure of a quadrantal angle is always a multiple of 90° , or $\frac{\pi}{2}$ radians.

EXAMPLE 2 Using the Unit Circle


Use the unit circle to evaluate the six trigonometric functions of $\theta = 270^\circ$.

SOLUTION

Step 1 Draw a unit circle with the angle $\theta = 270^\circ$ in standard position.

Step 2 Identify the point where the terminal side of θ intersects the unit circle. The terminal side of θ intersects the unit circle at $(0, -1)$.

Step 3 Find the values of the six trigonometric functions. Let $x = 0$ and $y = -1$ to evaluate the trigonometric functions.



$$\sin \theta = \frac{y}{r} = \frac{-1}{1} = -1$$

$$\csc \theta = \frac{r}{y} = \frac{1}{-1} = -1$$

$$\cos \theta = \frac{x}{r} = \frac{0}{1} = 0$$

$$\sec \theta = \frac{r}{x} = \frac{1}{0} \text{ undefined}$$

$$\tan \theta = \frac{y}{x} = \frac{-1}{0} \text{ undefined}$$




$$\cot \theta = \frac{x}{y} = \frac{0}{-1} = 0$$

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Evaluate the six trigonometric functions of θ .



4. Use the unit circle to evaluate the six trigonometric functions of $\theta = 180^\circ$.

Reference Angles


Core Concept

READING

The symbol θ' is read as "theta prime."

Reference Angle Relationships

Let θ be an angle in standard position. The **reference angle** for θ is the acute angle θ' formed by the terminal side of θ and the x -axis. The relationship between θ and θ' is shown below for nonquadrantal angles θ such that $90^\circ < \theta < 360^\circ$ or, in radians, $\frac{\pi}{2} < \theta < 2\pi$.



EXAMPLE 3 Finding Reference Angles

Find the reference angle θ' for (a) $\theta = \frac{5\pi}{3}$ and (b) $\theta = -130^\circ$.



SOLUTION

a. The terminal side of θ lies in Quadrant IV. So,

$$\theta' = 2\pi - \frac{5\pi}{3} = \frac{\pi}{3}.$$

$$\theta = \frac{5\pi}{3} \text{ and } \theta' = \frac{\pi}{3}.$$

b. Note that θ is coterminal with 230° , whose terminal side lies in Quadrant III. So, $\theta' = 230^\circ - 180^\circ = 50^\circ$. The figure at the left shows $\theta = -130^\circ$ and $\theta' = 50^\circ$.



Reference angles allow you to evaluate a trigonometric function for any angle θ . The sign of the trigonometric function value depends on the quadrant in which θ lies.

Core Concept

Evaluating Trigonometric Functions

Use these steps to evaluate a trigonometric function for any angle θ :

- Step 1** Find the reference angle θ' .
- Step 2** Evaluate the trigonometric function for θ' .
- Step 3** Determine the sign of the trigonometric function value from the quadrant in which θ lies.

Signs of Function Values

Quadrant II	Quadrant I
$\sin \theta, \csc \theta : +$	$\sin \theta, \csc \theta : +$
$\cos \theta, \sec \theta : -$	$\cos \theta, \sec \theta : +$
$\tan \theta, \cot \theta : -$	$\tan \theta, \cot \theta : +$
Quadrant III	Quadrant IV
$\sin \theta, \csc \theta : -$	$\sin \theta, \csc \theta : -$
$\cos \theta, \sec \theta : -$	$\cos \theta, \sec \theta : +$
$\tan \theta, \cot \theta : +$	$\tan \theta, \cot \theta : -$

EXAMPLE 4**Using Reference Angles to Evaluate Functions**

Evaluate (a) $\tan(-240^\circ)$ and (b) $\csc \frac{17\pi}{6}$.

SOLUTION

- a. The angle -240° is coterminal with 120° . The reference angle is $\theta' = 180^\circ - 120^\circ = 60^\circ$. The tangent function is negative in Quadrant II, so


$$\tan(-240^\circ) = -\tan 60^\circ = -\sqrt{3}.$$

- b. The angle $\frac{17\pi}{6}$ is coterminal with $\frac{5\pi}{6}$. The reference angle is

$$\theta' = \pi - \frac{5\pi}{6} = \frac{\pi}{6}.$$

The cosecant function is positive in Quadrant II, so

$$\csc \frac{17\pi}{6} = \csc \frac{\pi}{6} = 2.$$

**INTERPRETING MODELS**


This model neglects air resistance and assumes that the projectile's starting and ending heights are the same.

EXAMPLE 5**Solving a Real-Life Problem**

The horizontal distance d (in feet) traveled by a projectile launched at an angle θ and with an initial speed v (in feet per second) is given by

$$d = \frac{v^2}{32} \sin 2\theta. \quad \text{Model for horizontal distance}$$

Estimate the horizontal distance traveled by a golf ball that is hit at an angle of 50° with an initial speed of 105 feet per second.

**SOLUTION**

Note that the golf ball is launched at an angle of $\theta = 50^\circ$ with initial speed of $v = 105$ feet per second.

$$\begin{aligned} d &= \frac{v^2}{32} \sin 2\theta && \text{Write model for horizontal distance.} \\ &= \frac{105^2}{32} \sin(2 \cdot 50^\circ) && \text{Substitute 105 for } v \text{ and } 50^\circ \text{ for } \theta. \\ &\approx 339 && \text{Use a calculator.} \end{aligned}$$

► The golf ball travels a horizontal distance of about 339 feet.

Monitoring Progress

Help in English and Spanish at BigIdeasMath.com

Sketch the angle. Then find its reference angle.

5. 210°

6. -260°

7. $\frac{-7\pi}{9}$

8. $\frac{15\pi}{4}$

Evaluate the function without using a calculator.

9. $\cos(-210^\circ)$

10. $\sec \frac{11\pi}{4}$

11. Use the model given in Example 5 to estimate the horizontal distance traveled by a track and field long jumper who jumps at an angle of 20° and with an initial speed of 27 feet per second.

9.3 Exercises


Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** A(n) _____ is an angle in standard position whose terminal side lies on an axis.
- WRITING** Given an angle θ in standard position with its terminal side in Quadrant III, explain how you can use a reference angle to find $\cos \theta$.

Monitoring Progress and Modeling with Mathematics

In Exercises 3–8, evaluate the six trigonometric functions of θ . (See Example 1.)



In Exercises 9–14, use the unit circle to evaluate the six trigonometric functions of θ . (See Example 2.)

- $\theta = 0^\circ$
- $\theta = 540^\circ$
- $\theta = \frac{\pi}{2}$
- $\theta = \frac{7\pi}{2}$
- $\theta = -270^\circ$
- $\theta = -2\pi$

In Exercises 15–22, sketch the angle. Then find its reference angle. (See Example 3.)

- -100°
- 150°
- 320°
- -370°
- $\frac{15\pi}{4}$
- $\frac{8\pi}{3}$
- $-\frac{5\pi}{6}$
- $-\frac{13\pi}{6}$

23. **ERROR ANALYSIS** Let $(-3, 2)$ be a point on the terminal side of an angle θ in standard position. Describe and correct the error in finding $\tan \theta$.



$$\tan \theta = \frac{x}{y} = -\frac{3}{2}$$

24. **ERROR ANALYSIS** Describe and correct the error in finding a reference angle θ' for $\theta = 650^\circ$.



θ is coterminal with 290° , whose terminal side lies in Quadrant IV.

$$\text{So, } \theta' = 290^\circ - 270^\circ = 20^\circ.$$

In Exercises 25–32, evaluate the function without using a calculator. (See Example 4.)

- $\sec 135^\circ$
- $\tan 240^\circ$
- $\sin(-150^\circ)$
- $\csc(-420^\circ)$
- $\tan\left(-\frac{3\pi}{4}\right)$
- $\cot\left(\frac{-8\pi}{3}\right)$
- $\cos \frac{7\pi}{4}$
- $\sec \frac{11\pi}{6}$

In Exercises 33–36, use the model for horizontal distance given in Example 5.


33. You kick a football at an angle of 60° with an initial speed of 49 feet per second. Estimate the horizontal distance traveled by the football. (See Example 5.)
34. The “frogbot” is a robot designed for exploring rough terrain on other planets. It can jump at a 45° angle with an initial speed of 14 feet per second. Estimate the horizontal distance the frogbot can jump on Earth.



35. At what speed must the in-line skater launch himself off the ramp in order to land on the other side of the ramp?



36. To win a javelin throwing competition, your last throw must travel a horizontal distance of at least 100 feet. You release the javelin at a 40° angle with an initial speed of 71 feet per second. Do you win the competition? Justify your answer.
37. **MODELING WITH MATHEMATICS** A rock climber is using a rock climbing treadmill that is 10 feet long. The climber begins by lying horizontally on the treadmill, which is then rotated about its midpoint by 110° so that the rock climber is climbing toward the top. If the midpoint of the treadmill is 6 feet above the ground, how high above the ground is the top of the treadmill?



38. **REASONING** A Ferris wheel has a radius of 75 feet. You board a car at the bottom of the Ferris wheel, which is 10 feet above the ground, and rotate 255° counterclockwise before the ride temporarily stops. How high above the ground are you when the ride stops? If the radius of the Ferris wheel is doubled, is your height above the ground doubled? Explain your reasoning.


39. **DRAWING CONCLUSIONS** A sprinkler at ground level is used to water a garden. The water leaving the sprinkler has an initial speed of 25 feet per second.

- a. Use the model for horizontal distance given in Example 5 to complete the table.


Angle of sprinkler, θ	Horizontal distance water travels, d
30°	
35°	
40°	
45°	
50°	
55°	
60°	

- b. Which value of θ appears to maximize the horizontal distance traveled by the water? Use the model for horizontal distance and the unit circle to explain why your answer makes sense.
- c. Compare the horizontal distance traveled by the water when $\theta = (45 - k)^\circ$ with the distance when $\theta = (45 + k)^\circ$, for $0 < k < 45$.

40. **MODELING WITH MATHEMATICS** Your school’s marching band is performing at halftime during a football game. In the last formation, the band members form a circle 100 feet wide in the center of the field. You start at a point on the circle 100 feet from the goal line, march 300° around the circle, and then walk toward the goal line to exit the field. How far from the goal line are you at the point where you leave the circle?



41. **ANALYZING RELATIONSHIPS** Use symmetry and the given information to label the coordinates of the other points corresponding to special angles on the unit circle.




42. **THOUGHT PROVOKING** Use the interactive unit circle tool at *BigIdeasMath.com* to describe all values of θ for each situation.

- $\sin \theta > 0$, $\cos \theta < 0$, and $\tan \theta > 0$
- $\sin \theta > 0$, $\cos \theta < 0$, and $\tan \theta < 0$

43. **CRITICAL THINKING** Write $\tan \theta$ as the ratio of two other trigonometric functions. Use this ratio to explain why $\tan 90^\circ$ is undefined but $\cot 90^\circ = 0$.


44. **HOW DO YOU SEE IT?** Determine whether each of the six trigonometric functions of θ is *positive*, *negative*, or *zero*. Explain your reasoning.



45. **USING STRUCTURE** A line with slope m passes through the origin. An angle θ in standard position has a terminal side that coincides with the line. Use a trigonometric function to relate the slope of the line to the angle.


46. **MAKING AN ARGUMENT** Your friend claims that the only solution to the trigonometric equation $\tan \theta = \sqrt{3}$ is $\theta = 60^\circ$. Is your friend correct? Explain your reasoning.

47. **PROBLEM SOLVING** When two atoms in a molecule are bonded to a common atom, chemists are interested in both the bond angle and the lengths of the bonds. An ozone molecule is made up of two oxygen atoms bonded to a third oxygen atom, as shown.



- In the diagram, coordinates are given in picometers (pm). (Note: $1 \text{ pm} = 10^{-12} \text{ m}$) Find the coordinates (x, y) of the center of the oxygen atom in Quadrant II.
- Find the distance d (in picometers) between the centers of the two unbonded oxygen atoms.

48. **MATHEMATICAL CONNECTIONS** The latitude of a point on Earth is the degree measure of the shortest arc from that point to the equator. For example, the latitude of point P in the diagram equals the degree measure of arc PE . At what latitude θ is the circumference of the circle of latitude at P half the distance around the equator?



Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Find all real zeros of the polynomial function. (Section 4.6)

49. $f(x) = x^4 + 2x^3 + x^2 + 8x - 12$

50. $f(x) = x^5 + 4x^4 - 14x^3 - 14x^2 - 15x - 18$


Graph the function. (Section 4.8)

51. $f(x) = 2(x + 3)^2(x - 1)$

52. $f(x) = \frac{1}{3}(x - 4)(x + 5)(x + 9)$

53. $f(x) = x^2(x + 1)^3(x - 2)$

41. **ANALYZING RELATIONSHIPS** Use symmetry and the given information to label the coordinates of the other points corresponding to special angles on the unit circle.




42. **THOUGHT PROVOKING** Use the interactive unit circle tool at *BigIdeasMath.com* to describe all values of θ for each situation.

- $\sin \theta > 0$, $\cos \theta < 0$, and $\tan \theta > 0$
- $\sin \theta > 0$, $\cos \theta < 0$, and $\tan \theta < 0$

43. **CRITICAL THINKING** Write $\tan \theta$ as the ratio of two other trigonometric functions. Use this ratio to explain why $\tan 90^\circ$ is undefined but $\cot 90^\circ = 0$.


44. **HOW DO YOU SEE IT?** Determine whether each of the six trigonometric functions of θ is *positive*, *negative*, or *zero*. Explain your reasoning.




45. **USING STRUCTURE** A line with slope m passes through the origin. An angle θ in standard position has a terminal side that coincides with the line. Use a trigonometric function to relate the slope of the line to the angle.

46. **MAKING AN ARGUMENT** Your friend claims that the only solution to the trigonometric equation $\tan \theta = \sqrt{3}$ is $\theta = 60^\circ$. Is your friend correct? Explain your reasoning.

47. **PROBLEM SOLVING** When two atoms in a molecule are bonded to a common atom, chemists are interested in both the bond angle and the lengths of the bonds. An ozone molecule is made up of two oxygen atoms bonded to a third oxygen atom, as shown.



- In the diagram, coordinates are given in picometers (pm). (Note: $1 \text{ pm} = 10^{-12} \text{ m}$) Find the coordinates (x, y) of the center of the oxygen atom in Quadrant II.
 - Find the distance d (in picometers) between the centers of the two unbonded oxygen atoms.
48. **MATHEMATICAL CONNECTIONS** The latitude of a point on Earth is the degree measure of the shortest arc from that point to the equator. For example, the latitude of point P in the diagram equals the degree measure of arc PE . At what latitude θ is the circumference of the circle of latitude at P half the distance around the equator?



Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Find all real zeros of the polynomial function. (Section 4.6)

49. $f(x) = x^4 + 2x^3 + x^2 + 8x - 12$

50. $f(x) = x^5 + 4x^4 - 14x^3 - 14x^2 - 15x - 18$

Graph the function. (Section 4.8)

51. $f(x) = 2(x + 3)^2(x - 1)$

52. $f(x) = \frac{1}{3}(x - 4)(x + 5)(x + 9)$

53. $f(x) = x^2(x + 1)^3(x - 2)$

9.4 Lesson

Core Vocabulary

amplitude, p. 486
 periodic function, p. 486
 cycle, p. 486
 period, p. 486
 phase shift, p. 488
 midline, p. 488

Previous

transformations
 x-intercept


What You Will Learn

- ▶ Explore characteristics of sine and cosine functions.
- ▶ Stretch and shrink graphs of sine and cosine functions.
- ▶ Translate graphs of sine and cosine functions.
- ▶ Reflect graphs of sine and cosine functions.

Exploring Characteristics of Sine and Cosine Functions

In this lesson, you will learn to graph sine and cosine functions. The graphs of sine and cosine functions are related to the graphs of the parent functions $y = \sin x$ and $y = \cos x$, which are shown below.

x	-2π	$-\frac{3\pi}{2}$	$-\pi$	$-\frac{\pi}{2}$	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π
$y = \sin x$	0	1	0	-1	0	1	0	-1	0
$y = \cos x$	1	0	-1	0	1	0	-1	0	1



Core Concept

Characteristics of $y = \sin x$ and $y = \cos x$

- The domain of each function is all real numbers.
- The range of each function is $-1 \leq y \leq 1$. So, the minimum value of each function is -1 and the maximum value is 1 .
- The **amplitude** of the graph of each function is one-half of the difference of the maximum value and the minimum value, or $\frac{1}{2}[1 - (-1)] = 1$.
- Each function is **periodic**, which means that its graph has a repeating pattern. The shortest repeating portion of the graph is called a **cycle**. The horizontal length of each cycle is called the **period**. Each graph shown above has a period of 2π .
- The x -intercepts for $y = \sin x$ occur when $x = 0, \pm\pi, \pm2\pi, \pm3\pi, \dots$
- The x -intercepts for $y = \cos x$ occur when $x = \pm\frac{\pi}{2}, \pm\frac{3\pi}{2}, \pm\frac{5\pi}{2}, \pm\frac{7\pi}{2}, \dots$

Stretching and Shrinking Sine and Cosine Functions

The graphs of $y = a \sin bx$ and $y = a \cos bx$ represent transformations of their parent functions. The value of a indicates a vertical stretch ($a > 1$) or a vertical shrink ($0 < a < 1$) and changes the amplitude of the graph. The value of b indicates a horizontal stretch ($0 < b < 1$) or a horizontal shrink ($b > 1$) and changes the period of the graph.

REMEMBER

The graph of $y = a \cdot f(x)$ is a vertical stretch or shrink of the graph of $y = f(x)$ by a factor of a .

The graph of $y = f(bx)$ is a horizontal stretch or shrink of the graph of $y = f(x)$ by a factor of $\frac{1}{b}$.

$$y = a \sin bx$$

$$y = a \cos bx$$

vertical stretch or shrink by a factor of a horizontal stretch or shrink by a factor of $\frac{1}{b}$

Core Concept



Amplitude and Period

The amplitude and period of the graphs of $y = a \sin bx$ and $y = a \cos bx$, where a and b are nonzero real numbers, are as follows:

$$\text{Amplitude} = |a|$$

$$\text{Period} = \frac{2\pi}{|b|}$$

Each graph below shows five key points that partition the interval $0 \leq x \leq \frac{2\pi}{b}$ into four equal parts. You can use these points to sketch the graphs of $y = a \sin bx$ and $y = a \cos bx$. The **x-intercepts**, **maximum**, and **minimum** occur at these points.



EXAMPLE 1

Graphing a Sine Function

Identify the amplitude and period of $g(x) = 4 \sin x$. Then graph the function and describe the graph of g as a transformation of the graph of $f(x) = \sin x$.


SOLUTION

The function is of the form $g(x) = a \sin bx$ where $a = 4$ and $b = 1$. So, the amplitude is $a = 4$ and the period is $\frac{2\pi}{b} = \frac{2\pi}{1} = 2\pi$.

$$\text{Intercepts: } (0, 0); \left(\frac{1}{2} \cdot 2\pi, 0\right) = (\pi, 0); (2\pi, 0)$$


$$\text{Maximum: } \left(\frac{1}{4} \cdot 2\pi, 4\right) = \left(\frac{\pi}{2}, 4\right)$$

$$\text{Minimum: } \left(\frac{3}{4} \cdot 2\pi, -4\right) = \left(\frac{3\pi}{2}, -4\right)$$



REMEMBER

A vertical stretch of a graph does not change its x -intercept(s). So, it makes sense that the x -intercepts of $g(x) = 4 \sin x$ and $f(x) = \sin x$ are the same.



► The graph of g is a vertical stretch by a factor of 4 of the graph of f .

EXAMPLE 2 Graphing a Cosine Function

Identify the amplitude and period of $g(x) = \frac{1}{2} \cos 2\pi x$. Then graph the function and describe the graph of g as a transformation of the graph of $f(x) = \cos x$.


SOLUTION

The function is of the form $g(x) = a \cos bx$ where $a = \frac{1}{2}$ and $b = 2\pi$. So, the amplitude is $a = \frac{1}{2}$ and the period is $\frac{2\pi}{b} = \frac{2\pi}{2\pi} = 1$.

Intercepts: $\left(\frac{1}{4} \cdot 1, 0\right) = \left(\frac{1}{4}, 0\right); \left(\frac{3}{4} \cdot 1, 0\right) = \left(\frac{3}{4}, 0\right)$

Maximums: $\left(0, \frac{1}{2}\right); \left(1, \frac{1}{2}\right)$

Minimum: $\left(\frac{1}{2} \cdot 1, -\frac{1}{2}\right) = \left(\frac{1}{2}, -\frac{1}{2}\right)$



STUDY TIP

After you have drawn one complete cycle of the graph in Example 2 on the interval $0 \leq x \leq 1$, you can extend the graph by repeating the cycle as many times as desired to the left and right of $0 \leq x \leq 1$.

► The graph of g is a vertical shrink by a factor of $\frac{1}{2}$ and a horizontal shrink by a factor of $\frac{1}{2\pi}$ of the graph of f .

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Identify the amplitude and period of the function. Then graph the function and describe the graph of g as a transformation of the graph of its parent function.

1. $g(x) = \frac{1}{4} \sin x$ 2. $g(x) = \cos 2x$ 3. $g(x) = 2 \sin \pi x$ 4. $g(x) = \frac{1}{3} \cos \frac{1}{2}x$

REMEMBER

The graph of $y = f(x) + k$ is a vertical translation of the graph of $y = f(x)$.

The graph of $y = f(x - h)$ is a horizontal translation of the graph of $y = f(x)$.

Translating Sine and Cosine Functions

The graphs of $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$ represent translations of $y = a \sin bx$ and $y = a \cos bx$. The value of k indicates a translation up ($k > 0$) or down ($k < 0$). The value of h indicates a translation left ($h < 0$) or right ($h > 0$). A horizontal translation of a periodic function is called a **phase shift**.



Core Concept

Graphing $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$

To graph $y = a \sin b(x - h) + k$ or $y = a \cos b(x - h) + k$ where $a > 0$ and $b > 0$, follow these steps:

Step 1 Identify the amplitude a , the period $\frac{2\pi}{b}$, the horizontal shift h , and the vertical shift k of the graph.

Step 2 Draw the horizontal line $y = k$, called the **midline** of the graph.

Step 3 Find the five key points by translating the key points of $y = a \sin bx$ or $y = a \cos bx$ horizontally h units and vertically k units.

Step 4 Draw the graph through the five translated key points.

LOOKING FOR STRUCTURE

The graph of g is a translation 3 units up of the graph of $f(x) = 2 \sin 4x$. So, add 3 to the y -coordinates of the five key points of f .

EXAMPLE 3

Graphing a Vertical Translation

Graph $g(x) = 2 \sin 4x + 3$.

SOLUTION

Step 1 Identify the amplitude, period, horizontal shift, and vertical shift.

Amplitude: $a = 2$

Horizontal shift: $h = 0$

Period: $\frac{2\pi}{b} = \frac{2\pi}{4} = \frac{\pi}{2}$

Vertical shift: $k = 3$

Step 2 Draw the midline of the graph, $y = 3$.


Step 3 Find the five key points.

On $y = k$: $(0, 0 + 3) = (0, 3)$; $(\frac{\pi}{4}, 0 + 3) = (\frac{\pi}{4}, 3)$; $(\frac{\pi}{2}, 0 + 3) = (\frac{\pi}{2}, 3)$

Maximum: $(\frac{\pi}{8}, 2 + 3) = (\frac{\pi}{8}, 5)$

Minimum: $(\frac{3\pi}{8}, -2 + 3) = (\frac{3\pi}{8}, 1)$

Step 4 Draw the graph through the key points.



EXAMPLE 4

Graphing a Horizontal Translation

Graph $g(x) = 5 \cos \frac{1}{2}(x - 3\pi)$.

SOLUTION

Step 1 Identify the amplitude, period, horizontal shift, and vertical shift.

Amplitude: $a = 5$

Horizontal shift: $h = 3\pi$

Period: $\frac{2\pi}{b} = \frac{2\pi}{\frac{1}{2}} = 4\pi$

Vertical shift: $k = 0$

Step 2 Draw the midline of the graph. Because $k = 0$, the midline is the x -axis.


Step 3 Find the five key points.

On $y = k$: $(\pi + 3\pi, 0) = (4\pi, 0)$;
 $(3\pi + 3\pi, 0) = (6\pi, 0)$

Maximums: $(0 + 3\pi, 5) = (3\pi, 5)$;
 $(4\pi + 3\pi, 5) = (7\pi, 5)$

Minimum: $(2\pi + 3\pi, -5) = (5\pi, -5)$

Step 4 Draw the graph through the key points.



LOOKING FOR STRUCTURE

The graph of g is a translation 3π units right of the graph of $f(x) = 5 \cos \frac{1}{2}x$. So, add 3π to the x -coordinates of the five key points of f .

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Graph the function.

5. $g(x) = \cos x + 4$

6. $g(x) = \frac{1}{2} \sin \left(x - \frac{\pi}{2} \right)$



7. $g(x) = \sin(x + \pi) - 1$

Reflecting Sine and Cosine Functions

You have graphed functions of the form $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$, where $a > 0$ and $b > 0$. To see what happens when $a < 0$, consider the graphs of $y = -\sin x$ and $y = -\cos x$.

REMEMBER

This result makes sense because the graph of $y = -f(x)$ is a reflection in the x -axis of the graph of $y = f(x)$.



The graphs are reflections of the graphs of $y = \sin x$ and $y = \cos x$ in the x -axis. In general, when $a < 0$, the graphs of $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$ are reflections of the graphs of $y = |a| \sin b(x - h) + k$ and $y = |a| \cos b(x - h) + k$, respectively, in the midline $y = k$.

EXAMPLE 5 Graphing a Reflection

Graph $g(x) = -2 \sin \frac{2}{3} \left(x - \frac{\pi}{2} \right)$.

SOLUTION

Step 1 Identify the amplitude, period, horizontal shift, and vertical shift.

$$\text{Amplitude: } |a| = |-2| = 2$$

$$\text{Horizontal shift: } h = \frac{\pi}{2}$$

$$\text{Period: } \frac{2\pi}{b} = \frac{2\pi}{\frac{2}{3}} = 3\pi$$

$$\text{Vertical shift: } k = 0$$


Step 2 Draw the midline of the graph. Because $k = 0$, the midline is the x -axis.

Step 3 Find the five key points of $f(x) = |-2| \sin \frac{2}{3} \left(x - \frac{\pi}{2} \right)$.

$$\text{On } y = k: \left(0 + \frac{\pi}{2}, 0 \right) = \left(\frac{\pi}{2}, 0 \right); \left(\frac{3\pi}{2} + \frac{\pi}{2}, 0 \right) = (2\pi, 0); \left(3\pi + \frac{\pi}{2}, 0 \right) = \left(\frac{7\pi}{2}, 0 \right)$$

$$\text{Maximum: } \left(\frac{3\pi}{4} + \frac{\pi}{2}, 2 \right) = \left(\frac{5\pi}{4}, 2 \right) \quad \text{Minimum: } \left(\frac{9\pi}{4} + \frac{\pi}{2}, -2 \right) = \left(\frac{11\pi}{4}, -2 \right)$$

Step 4 Reflect the graph. Because $a < 0$, the graph is reflected in the midline $y = 0$. So, $\left(\frac{5\pi}{4}, 2 \right)$ becomes $\left(\frac{5\pi}{4}, -2 \right)$ and $\left(\frac{11\pi}{4}, -2 \right)$ becomes $\left(\frac{11\pi}{4}, 2 \right)$.



Step 5 Draw the graph through the key points.

STUDY TIP

In Example 5, the maximum value and minimum value of f are the minimum value and maximum value, respectively, of g .

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Graph the function.

$$8. g(x) = -\cos \left(x + \frac{\pi}{2} \right) \quad 9. g(x) = -3 \sin \frac{1}{2} x + 2 \quad 10. g(x) = -2 \cos 4x - 1$$

9.4 Exercises

Dynamic Solutions available at BigIdeasMath.com


Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** The shortest repeating portion of the graph of a periodic function is called a(n) _____.
- WRITING** Compare the amplitudes and periods of the functions $y = \frac{1}{2} \cos x$ and $y = 3 \cos 2x$.
- VOCABULARY** What is a phase shift? Give an example of a sine function that has a phase shift.
- VOCABULARY** What is the midline of the graph of the function $y = 2 \sin 3(x + 1) - 2$?


Monitoring Progress and Modeling with Mathematics

USING STRUCTURE In Exercises 5–8, determine whether the graph represents a periodic function. If so, identify the period.


5.




6.



7.




8.




In Exercises 9–12, identify the amplitude and period of the graph of the function.


9.




10.



11.



12.



In Exercises 13–20, identify the amplitude and period of the function. Then graph the function and describe the graph of g as a transformation of the graph of its parent function. (See Examples 1 and 2.)

13. $g(x) = 3 \sin x$

14. $g(x) = 2 \sin x$

15. $g(x) = \cos 3x$

16. $g(x) = \cos 4x$

17. $g(x) = \sin 2\pi x$

18. $g(x) = 3 \sin 2x$

19. $g(x) = \frac{1}{3} \cos 4x$

20. $g(x) = \frac{1}{2} \cos 4\pi x$

21. **ANALYZING EQUATIONS** Which functions have an amplitude of 4 and a period of 2?

(A) $y = 4 \cos 2x$

(B) $y = -4 \sin \pi x$

(C) $y = 2 \sin 4x$

(D) $y = 4 \cos \pi x$

22. **WRITING EQUATIONS** Write an equation of the form $y = a \sin bx$, where $a > 0$ and $b > 0$, so that the graph has the given amplitude and period.

a. amplitude: 1
period: 5

b. amplitude: 10
period: 4

c. amplitude: 2
period: 2π

d. amplitude: $\frac{1}{2}$
period: 3π

23. **MODELING WITH MATHEMATICS** The motion of a pendulum can be modeled by the function $d = 4 \cos 8\pi t$, where d is the horizontal displacement (in inches) of the pendulum relative to its position at rest and t is the time (in seconds). Find and interpret the period and amplitude in the context of this situation. Then graph the function.

- 24. MODELING WITH MATHEMATICS** A buoy bobs up and down as waves go past. The vertical displacement y (in feet) of the buoy with respect to sea level can be modeled by $y = 1.75 \cos \frac{\pi}{3}t$, where t is the time (in seconds). Find and interpret the period and amplitude in the context of the problem. Then graph the function.



In Exercises 25–34, graph the function. (See Examples 3 and 4.)

25. $g(x) = \sin x + 2$ 26. $g(x) = \cos x - 4$
 27. $g(x) = \cos\left(x - \frac{\pi}{2}\right)$ 28. $g(x) = \sin\left(x + \frac{\pi}{4}\right)$
 29. $g(x) = 2 \cos x - 1$ 30. $g(x) = 3 \sin x + 1$
 31. $g(x) = \sin 2(x + \pi)$
 32. $g(x) = \cos 2(x - \pi)$
 33. $g(x) = \sin \frac{1}{2}(x + 2\pi) + 3$
 34. $g(x) = \cos \frac{1}{2}(x - 3\pi) - 5$

- 35. ERROR ANALYSIS** Describe and correct the error in finding the period of the function $y = \sin \frac{2}{3}x$.



$$\text{Period: } \frac{|b|}{2\pi} = \frac{\left|\frac{2}{3}\right|}{2\pi} = \frac{1}{3\pi}$$

- 36. ERROR ANALYSIS** Describe and correct the error in determining the point where the maximum value of the function $y = 2 \sin\left(x - \frac{\pi}{2}\right)$ occurs.



Maximum:

$$\left(\left(\frac{1}{4} \cdot 2\pi\right) - \frac{\pi}{2}, 2\right) = \left(\frac{\pi}{2} - \frac{\pi}{2}, 2\right) = (0, 2)$$

USING STRUCTURE In Exercises 37–40, describe the transformation of the graph of f represented by the function g .

37. $f(x) = \cos x$, $g(x) = 2 \cos\left(x - \frac{\pi}{2}\right) + 1$
 38. $f(x) = \sin x$, $g(x) = 3 \sin\left(x + \frac{\pi}{4}\right) - 2$
 39. $f(x) = \sin x$, $g(x) = \sin 3(x + 3\pi) - 5$
 40. $f(x) = \cos x$, $g(x) = \cos 6(x - \pi) + 9$

In Exercises 41–48, graph the function. (See Example 5.)

41. $g(x) = -\cos x + 3$ 42. $g(x) = -\sin x - 5$
 43. $g(x) = -\sin \frac{1}{2}x - 2$ 44. $g(x) = -\cos 2x + 1$
 45. $g(x) = -\sin(x - \pi) + 4$
 46. $g(x) = -\cos(x + \pi) - 2$
 47. $g(x) = -4 \cos\left(x + \frac{\pi}{4}\right) - 1$
 48. $g(x) = -5 \sin\left(x - \frac{\pi}{2}\right) + 3$


- 49. USING EQUATIONS** Which of the following is a point where the maximum value of the graph of $y = -4 \cos\left(x - \frac{\pi}{2}\right)$ occurs?

- (A) $\left(-\frac{\pi}{2}, 4\right)$ (B) $\left(\frac{\pi}{2}, 4\right)$
 (C) $(0, 4)$ (D) $(\pi, 4)$


- 50. ANALYZING RELATIONSHIPS** Match each function with its graph. Explain your reasoning.

- a. $y = 3 + \sin x$ b. $y = -3 + \cos x$
 c. $y = \sin 2\left(x - \frac{\pi}{2}\right)$ d. $y = \cos 2\left(x - \frac{\pi}{2}\right)$


A.




B.



C.




D.



WRITING EQUATIONS In Exercises 51–54, write a rule for g that represents the indicated transformations of the graph of f .

51. $f(x) = 3 \sin x$; translation 2 units up and π units right
52. $f(x) = \cos 2\pi x$; translation 4 units down and 3 units left
53. $f(x) = \frac{1}{3} \cos \pi x$; translation 1 unit down, followed by a reflection in the line $y = -1$
54. $f(x) = \frac{1}{2} \sin 6x$; translation $\frac{3}{2}$ units down and 1 unit right, followed by a reflection in the line $y = -\frac{3}{2}$

55. **MODELING WITH MATHEMATICS** The height h (in feet) of a swing above the ground can be modeled by the function $h = -8 \cos \theta + 10$, where the pivot is 10 feet above the ground, the rope is 8 feet long, and θ is the angle that the rope makes with the vertical. Graph the function. What is the height of the swing when θ is 45° ?




56. **DRAWING A CONCLUSION** In a particular region, the population L (in thousands) of lynx (the predator) and the population H (in thousands) of hares (the prey) can be modeled by the equations

$$L = 11.5 + 6.5 \sin \frac{\pi}{5} t$$

$$H = 27.5 + 17.5 \cos \frac{\pi}{5} t$$

where t is the time in years.

- a. Determine the ratio of hares to lynx when $t = 0, 2.5, 5$, and 7.5 years.
- b. Use the figure to explain how the changes in the two populations appear to be related.




57. **USING TOOLS** The average wind speed s (in miles per hour) in the Boston Harbor can be approximated by

$$s = 3.38 \sin \frac{\pi}{180}(t + 3) + 11.6$$

where t is the time in days and $t = 0$ represents January 1. Use a graphing calculator to graph the function. On which days of the year is the average wind speed 10 miles per hour? Explain your reasoning.

58. **USING TOOLS** The water depth d (in feet) for the Bay of Fundy can be modeled by $d = 35 - 28 \cos \frac{\pi}{6.2} t$, where t is the time in hours and $t = 0$ represents midnight. Use a graphing calculator to graph the function. At what time(s) is the water depth 7 feet? Explain.




59. **MULTIPLE REPRESENTATIONS** Find the average rate of change of each function over the interval $0 < x < \pi$.

a. $y = 2 \cos x$

b.

x	0	$\frac{\pi}{2}$	π	$\frac{3\pi}{2}$	2π
$f(x) = -\cos x$	-1	0	1	0	-1



60. **REASONING** Consider the functions $y = \sin(-x)$ and $y = \cos(-x)$.

- a. Construct a table of values for each equation using the quadrantal angles in the interval $-2\pi \leq x \leq 2\pi$.
- b. Graph each function.
- c. Describe the transformations of the graphs of the parent functions.


- 61. MODELING WITH MATHEMATICS** You are riding a Ferris wheel that turns for 180 seconds. Your height h (in feet) above the ground at any time t (in seconds) can be modeled by the equation

$$h = 85 \sin \frac{\pi}{20}(t - 10) + 90.$$

- Graph the function.
- How many cycles does the Ferris wheel make in 180 seconds?
- What are your maximum and minimum heights?




- 62. HOW DO YOU SEE IT?** Use the graph to answer each question.



- Does the graph represent a function of the form $f(x) = a \sin bx$ or $f(x) = a \cos bx$? Explain.
 - Identify the maximum value, minimum value, period, and amplitude of the function.
- 63. FINDING A PATTERN** Write an expression in terms of the integer n that represents all the x -intercepts of the graph of the function $y = \cos 2x$. Justify your answer.
- 64. MAKING AN ARGUMENT** Your friend states that for functions of the form $y = a \sin bx$ and $y = a \cos bx$, the values of a and b affect the x -intercepts of the graph of the function. Is your friend correct? Explain.
- 65. CRITICAL THINKING** Describe a transformation of the graph of $f(x) = \sin x$ that results in the graph of $g(x) = \cos x$.

- 66. THOUGHT PROVOKING** Use a graphing calculator to find a function of the form $y = \sin b_1x + \cos b_2x$ whose graph matches that shown below.



- 67. PROBLEM SOLVING** For a person at rest, the blood pressure P (in millimeters of mercury) at time t (in seconds) is given by the function

$$P = 100 - 20 \cos \frac{8\pi}{3}t.$$

Graph the function. One cycle is equivalent to one heartbeat. What is the pulse rate (in heartbeats per minute) of the person?



- 68. PROBLEM SOLVING** The motion of a spring can be modeled by $y = A \cos kt$, where y is the vertical displacement (in feet) of the spring relative to its position at rest, A is the initial displacement (in feet), k is a constant that measures the elasticity of the spring, and t is the time (in seconds).
- You have a spring whose motion can be modeled by the function $y = 0.2 \cos 6t$. Find the initial displacement and the period of the spring. Then graph the function.
 - When a damping force is applied to the spring, the motion of the spring can be modeled by the function $y = 0.2e^{-4.5t} \cos 4t$. Graph this function. What effect does damping have on the motion?

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Simplify the rational expression, if possible. (Section 7.3)

69. $\frac{x^2 + x - 6}{x + 3}$

70. $\frac{x^3 - 2x^2 - 24x}{x^2 - 2x - 24}$

71. $\frac{x^2 - 4x - 5}{x^2 + 4x - 5}$

72. $\frac{x^2 - 16}{x^2 + x - 20}$

Find the least common multiple of the expressions. (Section 7.4)

73. $2x, 2(x - 5)$

74. $x^2 - 4, x + 2$

75. $x^2 + 8x + 12, x + 6$

9.1–9.4 What Did You Learn?

Core Vocabulary

sine, *p.* 462
cosine, *p.* 462
tangent, *p.* 462
cosecant, *p.* 462
secant, *p.* 462
cotangent, *p.* 462
initial side, *p.* 470
terminal side, *p.* 470

standard position, *p.* 470
coterminal, *p.* 471
radian, *p.* 471
sector, *p.* 472
central angle, *p.* 472
unit circle, *p.* 479
quadrantal angle, *p.* 479
reference angle, *p.* 480

amplitude, *p.* 486
periodic function, *p.* 486
cycle, *p.* 486
period, *p.* 486
phase shift, *p.* 488
midline, *p.* 488

Core Concepts

Section 9.1

Right Triangle Definitions of Trigonometric Functions, *p.* 462
Trigonometric Values for Special Angles, *p.* 463

Section 9.2

Angles in Standard Position, *p.* 470
Converting Between Degrees and Radians, *p.* 471

Degree and Radian Measures of Special Angles, *p.* 472
Arc Length and Area of a Sector, *p.* 472

Section 9.3

General Definitions of Trigonometric Functions, *p.* 478
The Unit Circle, *p.* 479

Reference Angle Relationships, *p.* 480
Evaluating Trigonometric Functions, *p.* 480

Section 9.4

Characteristics of $y = \sin x$ and $y = \cos x$, *p.* 486
Amplitude and Period, *p.* 487
Graphing $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$, *p.* 488

Mathematical Practices

1. Make a conjecture about the horizontal distances traveled in part (c) of Exercise 39 on page 483.
2. Explain why the quantities in part (a) of Exercise 56 on page 493 make sense in the context of the situation.

Study Skills

Form a Final Exam Study Group




Form a study group several weeks before the final exam. The intent of this group is to review what you have already learned while continuing to learn new material.



9.1–9.4 Quiz

1. In a right triangle, θ is an acute angle and $\sin \theta = \frac{2}{7}$. Evaluate the other five trigonometric functions of θ . (Section 9.1)

Find the value of x for the right triangle. (Section 9.1)



Draw an angle with the given measure in standard position. Then find one positive angle and one negative angle that are coterminal with the given angle. (Section 9.2)

5. 40°

6. $\frac{5\pi}{6}$

7. -960°




Convert the degree measure to radians or the radian measure to degrees. (Section 9.2)

8. $\frac{3\pi}{10}$


9. -60°

10. 72°

Evaluate the six trigonometric functions of θ . (Section 9.3)



14. Identify the amplitude and period of $g(x) = 3 \sin x$. Then graph the function and describe the graph of g as a transformation of the graph of $f(x) = \sin x$. (Section 9.4)
15. Identify the amplitude and period of $g(x) = \cos 5\pi x + 3$. Then graph the function and describe the graph of g as a transformation of the graph of $f(x) = \cos x$. (Section 9.4)
16. You are flying a kite at an angle of 70° . You have let out a total of 400 feet of string and are holding the reel steady 4 feet above the ground. (Section 9.1)
- How high above the ground is the kite?
 - A friend watching the kite estimates that the angle of elevation to the kite is 85° . How far from your friend are you standing?
17. The top of the Space Needle in Seattle, Washington, is a revolving, circular restaurant. The restaurant has a radius of 47.25 feet and makes one complete revolution in about an hour. You have dinner at a window table from 7:00 P.M. to 8:55 P.M. Compare the distance you revolve with the distance of a person seated 5 feet away from the windows. (Section 9.2)



9.5 Graphing Other Trigonometric Functions

Essential Question What are the characteristics of the graph of the tangent function?


EXPLORATION 1 Graphing the Tangent Function

Work with a partner.

- a. Complete the table for $y = \tan x$, where x is an angle measure in radians.

x	$-\frac{\pi}{2}$	$-\frac{\pi}{3}$	$-\frac{\pi}{4}$	$-\frac{\pi}{6}$	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
$y = \tan x$									
x	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{5\pi}{4}$	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$
$y = \tan x$									

- b. The graph of $y = \tan x$ has vertical asymptotes at x -values where $\tan x$ is undefined. Plot the points (x, y) from part (a). Then use the asymptotes to sketch the graph of $y = \tan x$.



MAKING SENSE OF PROBLEMS

To be proficient in math, you need to consider analogous problems and try special cases of the original problem in order to gain insight into its solution.

- c. For the graph of $y = \tan x$, identify the asymptotes, the x -intercepts, and the intervals for which the function is increasing or decreasing over $-\frac{\pi}{2} \leq x \leq \frac{3\pi}{2}$. Is the tangent function *even*, *odd*, or *neither*?

Communicate Your Answer

- What are the characteristics of the graph of the tangent function?
- Describe the asymptotes of the graph of $y = \cot x$ on the interval $-\frac{\pi}{2} < x < \frac{3\pi}{2}$.

9.5 Lesson

Core Vocabulary

Previous

asymptote
period
amplitude
x-intercept
transformations

What You Will Learn

- ▶ Explore characteristics of tangent and cotangent functions.
- ▶ Graph tangent and cotangent functions.
- ▶ Graph secant and cosecant functions.

Exploring Tangent and Cotangent Functions


The graphs of tangent and cotangent functions are related to the graphs of the parent functions $y = \tan x$ and $y = \cot x$, which are graphed below.

	$\leftarrow x \text{ approaches } -\frac{\pi}{2}$					$x \text{ approaches } \frac{\pi}{2} \rightarrow$			
x	$-\frac{\pi}{2}$	-1.57	-1.5	$-\frac{\pi}{4}$	0	$\frac{\pi}{4}$	1.5	1.57	$\frac{\pi}{2}$
$y = \tan x$	Undef.	-1256	-14.10	-1	0	1	14.10	1256	Undef.
	$\leftarrow \tan x \text{ approaches } -\infty$					$\tan x \text{ approaches } \infty \rightarrow$			

Because $\tan x = \frac{\sin x}{\cos x}$, $\tan x$ is undefined for x -values at which $\cos x = 0$, such as $x = \pm \frac{\pi}{2} \approx \pm 1.571$.

The table indicates that the graph has asymptotes at these values. The table represents one cycle of the graph, so the period of the graph is π .

You can use a similar approach to graph $y = \cot x$. Because $\cot x = \frac{\cos x}{\sin x}$, $\cot x$ is undefined for x -values at which $\sin x = 0$, which are multiples of π . The graph has asymptotes at these values. The period of the graph is also π .



Core Concept

Characteristics of $y = \tan x$ and $y = \cot x$

The functions $y = \tan x$ and $y = \cot x$ have the following characteristics.

- The domain of $y = \tan x$ is all real numbers except odd multiples of $\frac{\pi}{2}$. At these x -values, the graph has vertical asymptotes.
- The domain of $y = \cot x$ is all real numbers except multiples of π . At these x -values, the graph has vertical asymptotes.
- The range of each function is all real numbers. So, the functions do not have maximum or minimum values, and the graphs do not have an amplitude.
- The period of each graph is π .
- The x -intercepts for $y = \tan x$ occur when $x = 0, \pm\pi, \pm2\pi, \pm3\pi, \dots$
- The x -intercepts for $y = \cot x$ occur when $x = \pm\frac{\pi}{2}, \pm\frac{3\pi}{2}, \pm\frac{5\pi}{2}, \pm\frac{7\pi}{2}, \dots$

STUDY TIP

Odd multiples of $\frac{\pi}{2}$ are values such as these:

$$\begin{aligned} \pm 1 \cdot \frac{\pi}{2} &= \pm \frac{\pi}{2} \\ \pm 3 \cdot \frac{\pi}{2} &= \pm \frac{3\pi}{2} \\ \pm 5 \cdot \frac{\pi}{2} &= \pm \frac{5\pi}{2} \end{aligned}$$

Graphing Tangent and Cotangent Functions

The graphs of $y = a \tan bx$ and $y = a \cot bx$ represent transformations of their parent functions. The value of a indicates a vertical stretch ($a > 1$) or a vertical shrink ($0 < a < 1$). The value of b indicates a horizontal stretch ($0 < b < 1$) or a horizontal shrink ($b > 1$) and changes the period of the graph.


Core Concept

Period and Vertical Asymptotes of $y = a \tan bx$ and $y = a \cot bx$

The period and vertical asymptotes of the graphs of $y = a \tan bx$ and $y = a \cot bx$, where a and b are nonzero real numbers, are as follows.

- The period of the graph of each function is $\frac{\pi}{|b|}$.
- The vertical asymptotes for $y = a \tan bx$ are at odd multiples of $\frac{\pi}{2|b|}$.
- The vertical asymptotes for $y = a \cot bx$ are at multiples of $\frac{\pi}{|b|}$.


Each graph below shows five key x -values that you can use to sketch the graphs of $y = a \tan bx$ and $y = a \cot bx$ for $a > 0$ and $b > 0$. These are the **x-intercept**, the **x-values where the asymptotes occur**, and the **x-values halfway between the x-intercept and the asymptotes**. At each halfway point, the value of the function is either a or $-a$.



EXAMPLE 1

Graphing a Tangent Function

Graph one period of $g(x) = 2 \tan 3x$. Describe the graph of g as a transformation of the graph of $f(x) = \tan x$.



SOLUTION

The function is of the form $g(x) = a \tan bx$ where $a = 2$ and $b = 3$. So, the period is $\frac{\pi}{|b|} = \frac{\pi}{3}$.

Intercept: $(0, 0)$

Asymptotes: $x = \frac{\pi}{2|b|} = \frac{\pi}{2(3)}$, or $x = \frac{\pi}{6}$; $x = -\frac{\pi}{2|b|} = -\frac{\pi}{2(3)}$, or $x = -\frac{\pi}{6}$

Halfway points: $\left(\frac{\pi}{4b}, a\right) = \left(\frac{\pi}{4(3)}, 2\right) = \left(\frac{\pi}{12}, 2\right)$;

$\left(-\frac{\pi}{4b}, -a\right) = \left(-\frac{\pi}{4(3)}, -2\right) = \left(-\frac{\pi}{12}, -2\right)$

► The graph of g is a vertical stretch by a factor of 2 and a horizontal shrink by a factor of $\frac{1}{3}$ of the graph of f .

EXAMPLE 2 Graphing a Cotangent Function

Graph one period of $g(x) = \cot \frac{1}{2}x$. Describe the graph of g as a transformation of the graph of $f(x) = \cot x$.

SOLUTION

The function is of the form $g(x) = a \cot bx$ where $a = 1$ and $b = \frac{1}{2}$. So, the period is $\frac{\pi}{|b|} = \frac{\pi}{\frac{1}{2}} = 2\pi$.

Intercept: $\left(\frac{\pi}{2b}, 0\right) = \left(\frac{\pi}{2(\frac{1}{2})}, 0\right) = (\pi, 0)$

Asymptotes: $x = 0$; $x = \frac{\pi}{|b|} = \frac{\pi}{\frac{1}{2}}$, or $x = 2\pi$

Halfway points: $\left(\frac{\pi}{4b}, a\right) = \left(\frac{\pi}{4(\frac{1}{2})}, 1\right) = \left(\frac{\pi}{2}, 1\right)$; $\left(\frac{3\pi}{4b}, -a\right) = \left(\frac{3\pi}{4(\frac{1}{2})}, -1\right) = \left(\frac{3\pi}{2}, -1\right)$

► The graph of g is a horizontal stretch by a factor of 2 of the graph of f .

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

Graph one period of the function. Describe the graph of g as a transformation of the graph of its parent function.


- $g(x) = \tan 2x$
- $g(x) = \frac{1}{3} \cot x$
- $g(x) = 2 \cot 4x$
- $g(x) = 5 \tan \pi x$

STUDY TIP

Because $\sec x = \frac{1}{\cos x}$, $\sec x$ is undefined for x -values at which $\cos x = 0$. The graph of $y = \sec x$ has vertical asymptotes at these x -values. You can use similar reasoning to understand the vertical asymptotes of the graph of $y = \csc x$.

Graphing Secant and Cosecant Functions

The graphs of secant and cosecant functions are related to the graphs of the parent functions $y = \sec x$ and $y = \csc x$, which are shown below.



Core Concept

Characteristics of $y = \sec x$ and $y = \csc x$

The functions $y = \sec x$ and $y = \csc x$ have the following characteristics.

- The domain of $y = \sec x$ is all real numbers except odd multiples of $\frac{\pi}{2}$. At these x -values, the graph has vertical asymptotes.
- The domain of $y = \csc x$ is all real numbers except multiples of π . At these x -values, the graph has vertical asymptotes.
- The range of each function is $y \leq -1$ and $y \geq 1$. So, the graphs do not have an amplitude.
- The period of each graph is 2π .

To graph $y = a \sec bx$ or $y = a \csc bx$, first graph the function $y = a \cos bx$ or $y = a \sin bx$, respectively. Then use the asymptotes and several points to sketch a graph of the function. Notice that the value of b represents a horizontal stretch or shrink by a factor of $\frac{1}{b}$, so the period of $y = a \sec bx$ and $y = a \csc bx$ is $\frac{2\pi}{|b|}$.

EXAMPLE 3

Graphing a Secant Function

Graph one period of $g(x) = 2 \sec x$. Describe the graph of g as a transformation of the graph of $f(x) = \sec x$.


SOLUTION

Step 1 Graph the function $y = 2 \cos x$.

The period is $\frac{2\pi}{1} = 2\pi$.

Step 2 Graph asymptotes of g . Because the asymptotes of g occur when $2 \cos x = 0$, graph $x = -\frac{\pi}{2}$, $x = \frac{\pi}{2}$, and $x = \frac{3\pi}{2}$.

Step 3 Plot points on g , such as $(0, 2)$ and $(\pi, -2)$. Then use the asymptotes to sketch the curve.



► The graph of g is a vertical stretch by a factor of 2 of the graph of f .

EXAMPLE 4

Graphing a Cosecant Function


Graph one period of $g(x) = \frac{1}{2} \csc \pi x$. Describe the graph of g as a transformation of the graph of $f(x) = \csc x$.

SOLUTION

Step 1 Graph the function $y = \frac{1}{2} \sin \pi x$. The period is $\frac{2\pi}{\pi} = 2$.

Step 2 Graph asymptotes of g . Because the asymptotes of g occur when $\frac{1}{2} \sin \pi x = 0$, graph $x = 0$, $x = 1$, and $x = 2$.

Step 3 Plot points on g , such as $(\frac{1}{2}, \frac{1}{2})$ and $(\frac{3}{2}, -\frac{1}{2})$. Then use the asymptotes to sketch the curve.



► The graph of g is a vertical shrink by a factor of $\frac{1}{2}$ and a horizontal shrink by a factor of $\frac{1}{\pi}$ of the graph of f .

LOOKING FOR A PATTERN

In Examples 3 and 4, notice that the plotted points are on both graphs. Also, these points represent a local maximum on one graph and a local minimum on the other graph.

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Graph one period of the function. Describe the graph of g as a transformation of the graph of its parent function.

5. $g(x) = \csc 3x$ 6. $g(x) = \frac{1}{2} \sec x$ 7. $g(x) = 2 \csc 2x$ 8. $g(x) = 2 \sec \pi x$

9.5 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- WRITING** Explain why the graphs of the tangent, cotangent, secant, and cosecant functions do not have an amplitude.
- COMPLETE THE SENTENCE** The _____ and _____ functions are undefined for x -values at which $\sin x = 0$.
- COMPLETE THE SENTENCE** The period of the function $y = \sec x$ is _____, and the period of $y = \cot x$ is _____.
- WRITING** Explain how to graph a function of the form $y = a \sec bx$.

Monitoring Progress and Modeling with Mathematics

In Exercises 5–12, graph one period of the function. Describe the graph of g as a transformation of the graph of its parent function. (See Examples 1 and 2.)

- $g(x) = 2 \tan x$
- $g(x) = 3 \tan x$
- $g(x) = \cot 3x$
- $g(x) = \cot 2x$
- $g(x) = 3 \cot \frac{1}{4}x$
- $g(x) = 4 \cot \frac{1}{2}x$
- $g(x) = \frac{1}{2} \tan \pi x$
- $g(x) = \frac{1}{3} \tan 2\pi x$

- ERROR ANALYSIS** Describe and correct the error in finding the period of the function $y = \cot 3x$.



Period: $\frac{2\pi}{|b|} = \frac{2\pi}{3}$

- ERROR ANALYSIS** Describe and correct the error in describing the transformation of $f(x) = \tan x$ represented by $g(x) = 2 \tan 5x$.





A vertical stretch by a factor of 5 and a horizontal shrink by a factor of $\frac{1}{2}$.

- ANALYZING RELATIONSHIPS** Use the given graph to graph each function.

a. $f(x) = 3 \sec 2x$

b. $f(x) = 4 \csc 3x$



- USING EQUATIONS** Which of the following are asymptotes of the graph of $y = 3 \tan 4x$?

(A) $x = \frac{\pi}{8}$

(B) $x = \frac{\pi}{4}$

(C) $x = 0$


(D) $x = -\frac{5\pi}{8}$

In Exercises 17–24, graph one period of the function. Describe the graph of g as a transformation of the graph of its parent function. (See Examples 3 and 4.)


- $g(x) = 3 \csc x$
- $g(x) = 2 \csc x$
- $g(x) = \sec 4x$
- $g(x) = \sec 3x$
- $g(x) = \frac{1}{2} \sec \pi x$
- $g(x) = \frac{1}{4} \sec 2\pi x$
- $g(x) = \csc \frac{\pi}{2}x$
- $g(x) = \csc \frac{\pi}{4}x$

ATTENDING TO PRECISION In Exercises 25–28, use the graph to write a function of the form $y = a \tan bx$.


25.




26.



27.



28.



USING STRUCTURE In Exercises 29–34, match the equation with the correct graph. Explain your reasoning.

29. $g(x) = 4 \tan x$

30. $g(x) = 4 \cot x$


31. $g(x) = 4 \csc \pi x$

32. $g(x) = 4 \sec \pi x$


33. $g(x) = \sec 2x$

34. $g(x) = \csc 2x$


A.




B.




C.




D.



E.



F.



35. **WRITING** Explain why there is more than one tangent function whose graph passes through the origin and has asymptotes at $x = -\pi$ and $x = \pi$.

36. **USING EQUATIONS** Graph one period of each function. Describe the transformation of the graph of its parent function.

a. $g(x) = \sec x + 3$

b. $g(x) = \csc x - 2$

c. $g(x) = \cot(x - \pi)$

d. $g(x) = -\tan x$

WRITING EQUATIONS In Exercises 37–40, write a rule for g that represents the indicated transformation of the graph of f .

37. $f(x) = \cot 2x$; translation 3 units up and $\frac{\pi}{2}$ units left

38. $f(x) = 2 \tan x$; translation π units right, followed by a horizontal shrink by a factor of $\frac{1}{3}$


39. $f(x) = 5 \sec(x - \pi)$; translation 2 units down, followed by a reflection in the x -axis

40. $f(x) = 4 \csc x$; vertical stretch by a factor of 2 and a reflection in the x -axis

41. **MULTIPLE REPRESENTATIONS** Which function has a greater local maximum value? Which has a greater local minimum value? Explain.


A. $f(x) = \frac{1}{4} \csc \pi x$

B.




42. **ANALYZING RELATIONSHIPS** Order the functions from the least average rate of change to the greatest average rate of change over the interval $-\frac{\pi}{4} < x < \frac{\pi}{4}$.


A.




B.




C.



D.




43. **REASONING** You are standing on a bridge 140 feet above the ground. You look down at a car traveling away from the underpass. The distance d (in feet) the car is from the base of the bridge can be modeled by $d = 140 \tan \theta$. Graph the function. Describe what happens to θ as d increases.



44. **USING TOOLS** You use a video camera to pan up the Statue of Liberty. The height h (in feet) of the part of the Statue of Liberty that can be seen through your video camera after time t (in seconds) can be modeled by $h = 100 \tan \frac{\pi}{36} t$. Graph the function using a graphing calculator. What viewing window did you use? Explain.

45. **MODELING WITH MATHEMATICS** You are standing 120 feet from the base of a 260-foot building. You watch your friend go down the side of the building in a glass elevator.



- Write an equation that gives the distance d (in feet) your friend is from the top of the building as a function of the angle of elevation θ .
- Graph the function found in part (a). Explain how the graph relates to this situation.


46. **MODELING WITH MATHEMATICS** You are standing 300 feet from the base of a 200-foot cliff. Your friend is rappelling down the cliff.

- Write an equation that gives the distance d (in feet) your friend is from the top of the cliff as a function of the angle of elevation θ .
- Graph the function found in part (a).
- Use a graphing calculator to determine the angle of elevation when your friend has rappelled halfway down the cliff.



47. **MAKING AN ARGUMENT** Your friend states that it is not possible to write a cosecant function that has the same graph as $y = \sec x$. Is your friend correct? Explain your reasoning.

48. **HOW DO YOU SEE IT?** Use the graph to answer each question.



- What is the period of the graph?
 - What is the range of the function?
 - Is the function of the form $f(x) = a \csc bx$ or $f(x) = a \sec bx$? Explain.
49. **ABSTRACT REASONING** Rewrite $a \sec bx$ in terms of $\cos bx$. Use your results to explain the relationship between the local maximums and minimums of the cosine and secant functions.

50. **THOUGHT PROVOKING** A trigonometric equation that is true for all values of the variable for which both sides of the equation are defined is called a *trigonometric identity*. Use a graphing calculator to graph the function

$$y = \frac{1}{2} \left(\tan \frac{x}{2} + \cot \frac{x}{2} \right).$$

Use your graph to write a trigonometric identity involving this function. Explain your reasoning.

51. **CRITICAL THINKING** Find a tangent function whose graph intersects the graph of $y = 2 + 2 \sin x$ only at minimum points of the sine function.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Write a cubic function whose graph passes through the given points. (Section 4.9)

52. $(-1, 0), (1, 0), (3, 0), (0, 3)$


53. $(-2, 0), (1, 0), (3, 0), (0, -6)$

54. $(-1, 0), (2, 0), (3, 0), (1, -2)$


55. $(-3, 0), (-1, 0), (3, 0), (-2, 1)$

Find the amplitude and period of the graph of the function. (Section 9.4)


56.



57.



58.



9.6 Modeling with Trigonometric Functions

Essential Question What are the characteristics of the real-life problems that can be modeled by trigonometric functions?


EXPLORATION 1 Modeling Electric Currents

MODELING WITH MATHEMATICS


To be proficient in math, you need to apply the mathematics you know to solve problems arising in everyday life.

Work with a partner. Find a sine function that models the electric current shown in each oscilloscope screen. State the amplitude and period of the graph.


a.




b.




c.




d.



e.



f.



Communicate Your Answer

- What are the characteristics of the real-life problems that can be modeled by trigonometric functions?
- Use the Internet or some other reference to find examples of real-life situations that can be modeled by trigonometric functions.

9.6 Lesson

Core Vocabulary

frequency, p. 506

sinusoid, p. 507

Previous

amplitude

period

midline

What You Will Learn


- Interpret and use frequency.
- Write trigonometric functions.
- Use technology to find trigonometric models.

Frequency

The periodic nature of trigonometric functions makes them useful for modeling *oscillating* motions or repeating patterns that occur in real life. Some examples are sound waves, the motion of a pendulum, and seasons of the year. In such applications, the reciprocal of the period is called the **frequency**, which gives the number of cycles per unit of time.

EXAMPLE 1 Using Frequency

A sound consisting of a single frequency is called a *pure tone*. An audiometer produces pure tones to test a person's auditory functions. An audiometer produces a pure tone with a frequency f of 2000 hertz (cycles per second). The maximum pressure P produced from the pure tone is 2 millipascals. Write and graph a sine model that gives the pressure P as a function of the time t (in seconds).



SOLUTION

Step 1 Find the values of a and b in the model $P = a \sin bt$. The maximum pressure is 2, so $a = 2$. Use the frequency f to find b .

$$\text{frequency} = \frac{1}{\text{period}} \quad \text{Write relationship involving frequency and period.}$$

$$2000 = \frac{b}{2\pi} \quad \text{Substitute.}$$

$$4000\pi = b \quad \text{Multiply each side by } 2\pi.$$

The pressure P as a function of time t is given by $P = 2 \sin 4000\pi t$.

Step 2 Graph the model. The amplitude is $a = 2$ and the period is

$$\frac{1}{f} = \frac{1}{2000}.$$


The key points are:

$$\text{Intercepts: } (0, 0); \left(\frac{1}{2} \cdot \frac{1}{2000}, 0\right) = \left(\frac{1}{4000}, 0\right); \left(\frac{1}{2000}, 0\right)$$

$$\text{Maximum: } \left(\frac{1}{4} \cdot \frac{1}{2000}, 2\right) = \left(\frac{1}{8000}, 2\right)$$

$$\text{Minimum: } \left(\frac{3}{4} \cdot \frac{1}{2000}, -2\right) = \left(\frac{3}{8000}, -2\right)$$

► The graph of $P = 2 \sin 4000\pi t$ is shown at the left.



1. **WHAT IF?** In Example 1, how would the function change when the audiometer produced a pure tone with a frequency of 1000 hertz?

Writing Trigonometric Functions


Graphs of sine and cosine functions are called **sinusoids**. One method to write a sine or cosine function that models a sinusoid is to find the values of a , b , h , and k for

$$y = a \sin b(x - h) + k \quad \text{or} \quad y = a \cos b(x - h) + k$$

where $|a|$ is the amplitude, $\frac{2\pi}{b}$ is the period ($b > 0$), h is the horizontal shift, and k is the vertical shift.

EXAMPLE 2 Writing a Trigonometric Function

Write a function for the sinusoid shown.



SOLUTION

Step 1 Find the maximum and minimum values. From the graph, the maximum value is 5 and the minimum value is -1 .

Step 2 Identify the vertical shift, k . The value of k is the mean of the maximum and minimum values.

$$k = \frac{(\text{maximum value}) + (\text{minimum value})}{2} = \frac{5 + (-1)}{2} = \frac{4}{2} = 2$$

Step 3 Decide whether the graph should be modeled by a sine or cosine function. Because the graph crosses the midline $y = 2$ on the y -axis, the graph is a sine curve with no horizontal shift. So, $h = 0$.

Step 4 Find the amplitude and period. The period is

$$\frac{\pi}{2} = \frac{2\pi}{b} \quad \Rightarrow \quad b = 4.$$

The amplitude is

$$|a| = \frac{(\text{maximum value}) - (\text{minimum value})}{2} = \frac{5 - (-1)}{2} = \frac{6}{2} = 3.$$


The graph is not a reflection, so $a > 0$. Therefore, $a = 3$.

► The function is $y = 3 \sin 4x + 2$. Check this by graphing the function on a graphing calculator.

STUDY TIP


Because the graph repeats every $\frac{\pi}{2}$ units, the period is $\frac{\pi}{2}$.

Check



EXAMPLE 3 Modeling Circular Motion

Two people swing jump ropes, as shown in the diagram. The highest point of the middle of each rope is 75 inches above the ground, and the lowest point is 3 inches. The rope makes 2 revolutions per second. Write a model for the height h (in inches) of a rope as a function of the time t (in seconds) given that the rope is at its lowest point when $t = 0$.



SOLUTION

A rope oscillates between 3 inches and 75 inches above the ground. So, a sine or cosine function may be an appropriate model for the height over time.

Step 1 Identify the maximum and minimum values. The maximum height of a rope is 75 inches. The minimum height is 3 inches.

Step 2 Identify the vertical shift, k .

$$k = \frac{(\text{maximum value}) + (\text{minimum value})}{2} = \frac{75 + 3}{2} = 39$$

Step 3 Decide whether the height should be modeled by a sine or cosine function. When $t = 0$, the height is at its minimum. So, use a cosine function whose graph is a reflection in the x -axis with no horizontal shift ($h = 0$).

Step 4 Find the amplitude and period.

$$\text{The amplitude is } |a| = \frac{(\text{maximum value}) - (\text{minimum value})}{2} = \frac{75 - 3}{2} = 36.$$

Because the graph is a reflection in the x -axis, $a < 0$. So, $a = -36$. Because a rope is rotating at a rate of 2 revolutions per second, one revolution is completed in 0.5 second. So, the period is $\frac{2\pi}{b} = 0.5$, and $b = 4\pi$.

► A model for the height of a rope is $h(t) = -36 \cos 4\pi t + 39$.

Check

Use the *table* feature of a graphing calculator to check your model.

X	Y1
0	3
.25	75
.5	3
.75	75
1	3
1.25	75
1.5	3


$X=0$

2 revolutions


Monitoring Progress Help in English and Spanish at BigIdeasMath.com

Write a function for the sinusoid.

2.



3.



4. **WHAT IF?** Describe how the model in Example 3 changes when the lowest point of a rope is 5 inches above the ground and the highest point is 70 inches above the ground.

Using Technology to Find Trigonometric Models

Another way to model sinusoids is to use a graphing calculator that has a sinusoidal regression feature.

EXAMPLE 4 Using Sinusoidal Regression



The table shows the numbers N of hours of daylight in Denver, Colorado, on the 15th day of each month, where $t = 1$ represents January. Write a model that gives N as a function of t and interpret the period of its graph.

t	1	2	3	4	5	6
N	9.68	10.75	11.93	13.27	14.38	14.98


t	7	8	9	10	11	12
N	14.70	13.73	12.45	11.17	9.98	9.38

SOLUTION

Step 1 Enter the data in a graphing calculator.

L1	L2	L3	1
1	9.68		
2	10.75		
3	11.93		
4	13.27		
5	14.38		
6	14.98		
7	14.7		
L1(1)=1			


Step 2 Make a scatter plot.



Step 3 The scatter plot appears sinusoidal. So, perform a sinusoidal regression.

```
SinReg
y=a*sin(bx+c)+d
a=2.764734198
b=.5111635715
c=-1.591149599
d=12.13293913
```

Step 4 Graph the data and the model in the same viewing window.



STUDY TIP

Notice that the *sinusoidal regression* feature finds a model of the form $y = a \sin(bx + c) + d$. This function has a period of $\frac{2\pi}{b}$ because it can be written as $y = a \sin b\left(x + \frac{c}{b}\right) + d$.

► The model appears to be a good fit. So, a model for the data is $N = 2.76 \sin(0.511t - 1.59) + 12.1$. The period, $\frac{2\pi}{0.511} \approx 12$, makes sense because there are 12 months in a year and you would expect this pattern to continue in following years.

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

5. The table shows the average daily temperature T (in degrees Fahrenheit) for a city each month, where $m = 1$ represents January. Write a model that gives T as a function of m and interpret the period of its graph.


m	1	2	3	4	5	6	7	8	9	10	11	12
T	29	32	39	48	59	68	74	72	65	54	45	35

9.6 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** Graphs of sine and cosine functions are called _____.
- WRITING** Describe how to find the frequency of the function whose graph is shown.



Monitoring Progress and Modeling with Mathematics

In Exercises 3–10, find the frequency of the function.





- $y = \sin x$
- $y = \sin 3x$
- $y = \cos 4x + 2$
- $y = -\cos 2x$
- $y = \sin 3\pi x$
- $y = \cos \frac{\pi x}{4}$
- $y = \frac{1}{2} \cos 0.75x - 8$
- $y = 3 \sin 0.2x + 6$

- MODELING WITH MATHEMATICS** The lowest frequency of sounds that can be heard by humans is 20 hertz. The maximum pressure P produced from a sound with a frequency of 20 hertz is 0.02 millipascal. Write and graph a sine model that gives the pressure P as a function of the time t (in seconds). (See Example 1.)

- MODELING WITH MATHEMATICS** A middle-A tuning fork vibrates with a frequency f of 440 hertz (cycles per second). You strike a middle-A tuning fork with a force that produces a maximum pressure of 5 pascals. Write and graph a sine model that gives the pressure P as a function of the time t (in seconds).



In Exercises 13–16, write a function for the sinusoid.
(See Example 2.)



17. **ERROR ANALYSIS** Describe and correct the error in finding the amplitude of a sinusoid with a maximum point at $(2, 10)$ and a minimum point at $(4, -6)$.

X

$$|a| = \frac{(\text{maximum value}) + (\text{minimum value})}{2}$$

$$= \frac{10 - 6}{2}$$

$$= 2$$

18. **ERROR ANALYSIS** Describe and correct the error in finding the vertical shift of a sinusoid with a maximum point at $(3, -2)$ and a minimum point at $(7, -8)$.

X

$$k = \frac{(\text{maximum value}) + (\text{minimum value})}{2}$$

$$= \frac{7 + 3}{2}$$

$$= 5$$

19. **MODELING WITH MATHEMATICS** One of the largest sewing machines in the world has a *flywheel* (which turns as the machine sews) that is 5 feet in diameter. The highest point of the handle at the edge of the flywheel is 9 feet above the ground, and the lowest point is 4 feet. The wheel makes a complete turn every 2 seconds. Write a model for the height h (in feet) of the handle as a function of the time t (in seconds) given that the handle is at its lowest point when $t = 0$. (See Example 3.)

20. **MODELING WITH MATHEMATICS** The Great Laxey Wheel, located on the Isle of Man, is the largest working water wheel in the world. The highest point of a bucket on the wheel is 70.5 feet above the viewing platform, and the lowest point is 2 feet below the viewing platform. The wheel makes a complete turn every 24 seconds. Write a model for the height h (in feet) of the bucket as a function of time t (in seconds) given that the bucket is at its lowest point when $t = 0$.



USING TOOLS In Exercises 21 and 22, the time t is measured in months, where $t = 1$ represents January. Write a model that gives the average monthly high temperature D as a function of t and interpret the period of the graph. (See Example 4.)


21.

Air Temperatures in Apple Valley, CA						
t	1	2	3	4	5	6
D	60	63	69	75	85	94
t	7	8	9	10	11	12
D	99	99	93	81	69	60


22.

Water Temperatures at Miami Beach, FL						
t	1	2	3	4	5	6
D	71	73	75	78	81	85
t	7	8	9	10	11	12
D	86	85	84	81	76	73

23. **MODELING WITH MATHEMATICS** A circuit has an alternating voltage of 100 volts that peaks every 0.5 second. Write a sinusoidal model for the voltage V as a function of the time t (in seconds).



24. **MULTIPLE REPRESENTATIONS** The graph shows the average daily temperature of Lexington, Kentucky. The average daily temperature of Louisville, Kentucky, is modeled by $y = -22 \cos \frac{\pi}{6}t + 57$, where y is the temperature (in degrees Fahrenheit) and t is the number of months since January 1. Which city has the greater average daily temperature? Explain.




25. **USING TOOLS** The table shows the numbers of employees N (in thousands) at a sporting goods company each year for 11 years. The time t is measured in years, with $t = 1$ representing the first year.

t	1	2	3	4	5	6
N	20.8	22.7	24.6	23.2	20	17.5

t	7	8	9	10	11	12
N	16.7	17.8	21	22	24.1	


- Use sinusoidal regression to find a model that gives N as a function of t .
- Predict the number of employees at the company in the 12th year.

26. **THOUGHT PROVOKING** The figure shows a tangent line drawn to the graph of the function $y = \sin x$. At several points on the graph, draw a tangent line to the graph and estimate its slope. Then plot the points (x, m) , where m is the slope of the tangent line. What can you conclude?




27. **REASONING** Determine whether you would use a sine or cosine function to model each sinusoid with the y -intercept described. Explain your reasoning.
- The y -intercept occurs at the maximum value of the function.
 - The y -intercept occurs at the minimum value of the function.
 - The y -intercept occurs halfway between the maximum and minimum values of the function.

28. **HOW DO YOU SEE IT?** What is the frequency of the function whose graph is shown? Explain.



29. **USING STRUCTURE** During one cycle, a sinusoid has a minimum at $(\frac{\pi}{2}, 3)$ and a maximum at $(\frac{\pi}{4}, 8)$. Write a sine function *and* a cosine function for the sinusoid. Use a graphing calculator to verify that your answers are correct.
30. **MAKING AN ARGUMENT** Your friend claims that a function with a frequency of 2 has a greater period than a function with a frequency of $\frac{1}{2}$. Is your friend correct? Explain your reasoning.
31. **PROBLEM SOLVING** The low tide at a port is 3.5 feet and occurs at midnight. After 6 hours, the port is at high tide, which is 16.5 feet.



- Write a sinusoidal model that gives the tide depth d (in feet) as a function of the time t (in hours). Let $t = 0$ represent midnight.
- Find all the times when low and high tides occur in a 24-hour period.
- Explain how the graph of the function you wrote in part (a) is related to a graph that shows the tide depth d at the port t hours after 3:00 A.M.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Simplify the expression. (Section 5.2)

32. $\frac{17}{\sqrt{2}}$

33. $\frac{3}{\sqrt{6} - 2}$

34. $\frac{8}{\sqrt{10} + 3}$

35. $\frac{13}{\sqrt{3} + \sqrt{11}}$

Expand the logarithmic expression. (Section 6.5)

36. $\log_8 \frac{x}{7}$

37. $\ln 2x$

38. $\log_3 5x^3$


39. $\ln \frac{4x^6}{y}$

9.7 Using Trigonometric Identities

Essential Question How can you verify a trigonometric identity?

EXPLORATION 1 Writing a Trigonometric Identity

Work with a partner. In the figure, the point (x, y) is on a circle of radius c with center at the origin.



- Write an equation that relates a , b , and c .
- Write expressions for the sine and cosine ratios of angle θ .
- Use the results from parts (a) and (b) to find the sum of $\sin^2 \theta$ and $\cos^2 \theta$. What do you observe?
- Complete the table to verify that the identity you wrote in part (c) is valid for angles (of your choice) in each of the four quadrants.

	θ	$\sin^2 \theta$	$\cos^2 \theta$	$\sin^2 \theta + \cos^2 \theta$
QI				
QII				
QIII				
QIV				

REASONING ABSTRACTLY

To be proficient in math, you need to know and flexibly use different properties of operations and objects.

EXPLORATION 2 Writing Other Trigonometric Identities

Work with a partner. The trigonometric identity you derived in Exploration 1 is called a Pythagorean identity. There are two other Pythagorean identities. To derive them, recall the four relationships:

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta} \qquad \csc \theta = \frac{1}{\sin \theta}$$

- Divide each side of the Pythagorean identity you derived in Exploration 1 by $\cos^2 \theta$ and simplify. What do you observe?
- Divide each side of the Pythagorean identity you derived in Exploration 1 by $\sin^2 \theta$ and simplify. What do you observe?

Communicate Your Answer

- How can you verify a trigonometric identity?
- Is $\sin \theta = \cos \theta$ a trigonometric identity? Explain your reasoning.
- Give some examples of trigonometric identities that are different than those in Explorations 1 and 2.

9.7 Lesson

Core Vocabulary

trigonometric identity, p. 514

Previous

unit circle

STUDY TIP

Note that $\sin^2 \theta$ represents $(\sin \theta)^2$ and $\cos^2 \theta$ represents $(\cos \theta)^2$.

What You Will Learn

- ▶ Use trigonometric identities to evaluate trigonometric functions and simplify trigonometric expressions.
- ▶ Verify trigonometric identities.


Using Trigonometric Identities

Recall that when an angle θ is in standard position with its terminal side intersecting the unit circle at (x, y) , then $x = \cos \theta$ and $y = \sin \theta$. Because (x, y) is on a circle centered at the origin with radius 1, it follows that

$$x^2 + y^2 = 1$$

and

$$\cos^2 \theta + \sin^2 \theta = 1.$$



The equation $\cos^2 \theta + \sin^2 \theta = 1$ is true for any value of θ . A trigonometric equation that is true for all values of the variable for which both sides of the equation are defined is called a **trigonometric identity**. In Section 9.1, you used reciprocal identities to find the values of the cosecant, secant, and cotangent functions. These and other fundamental trigonometric identities are listed below.

Core Concept

Fundamental Trigonometric Identities

Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$

Tangent and Cotangent Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

Cofunction Identities

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta$$

Negative Angle Identities

$$\sin(-\theta) = -\sin \theta$$

$$\cos(-\theta) = \cos \theta$$

$$\tan(-\theta) = -\tan \theta$$

In this section, you will use trigonometric identities to do the following.

- Evaluate trigonometric functions.
- Simplify trigonometric expressions.
- Verify other trigonometric identities.

EXAMPLE 1**Finding Trigonometric Values**

Given that $\sin \theta = \frac{4}{5}$ and $\frac{\pi}{2} < \theta < \pi$, find the values of the other five trigonometric functions of θ .

SOLUTION

Step 1 Find $\cos \theta$.

$$\sin^2 \theta + \cos^2 \theta = 1$$

Write Pythagorean identity.

$$\left(\frac{4}{5}\right)^2 + \cos^2 \theta = 1$$

Substitute $\frac{4}{5}$ for $\sin \theta$.

$$\cos^2 \theta = 1 - \left(\frac{4}{5}\right)^2$$

Subtract $\left(\frac{4}{5}\right)^2$ from each side.

$$\cos^2 \theta = \frac{9}{25}$$

Simplify.

$$\cos \theta = \pm \frac{3}{5}$$

Take square root of each side.

$$\cos \theta = -\frac{3}{5}$$

Because θ is in Quadrant II, $\cos \theta$ is negative.

Step 2 Find the values of the other four trigonometric functions of θ using the values of $\sin \theta$ and $\cos \theta$.

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{4}{5}}{-\frac{3}{5}} = -\frac{4}{3} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{-\frac{3}{5}}{\frac{4}{5}} = -\frac{3}{4}$$

$$\csc \theta = \frac{1}{\sin \theta} = \frac{1}{\frac{4}{5}} = \frac{5}{4} \qquad \sec \theta = \frac{1}{\cos \theta} = \frac{1}{-\frac{3}{5}} = -\frac{5}{3}$$

EXAMPLE 2**Simplifying Trigonometric Expressions**

Simplify (a) $\tan\left(\frac{\pi}{2} - \theta\right)\sin \theta$ and (b) $\sec \theta \tan^2 \theta + \sec \theta$.

SOLUTION

a. $\tan\left(\frac{\pi}{2} - \theta\right)\sin \theta = \cot \theta \sin \theta$

Cofunction identity

$$= \left(\frac{\cos \theta}{\sin \theta}\right)(\sin \theta)$$

Cotangent identity

$$= \cos \theta$$

Simplify.

b. $\sec \theta \tan^2 \theta + \sec \theta = \sec \theta (\sec^2 \theta - 1) + \sec \theta$

Pythagorean identity

$$= \sec^3 \theta - \sec \theta + \sec \theta$$

Distributive Property

$$= \sec^3 \theta$$

Simplify.

Monitoring Progress

Help in English and Spanish at BigIdeasMath.com

1. Given that $\cos \theta = \frac{1}{6}$ and $0 < \theta < \frac{\pi}{2}$, find the values of the other five trigonometric functions of θ .

Simplify the expression.

2. $\sin x \cot x \sec x$

3. $\cos \theta - \cos \theta \sin^2 \theta$

4. $\frac{\tan x \csc x}{\sec x}$

Verifying Trigonometric Identities

You can use the fundamental identities from this chapter to verify new trigonometric identities. When verifying an identity, begin with the expression on one side. Use algebra and trigonometric properties to manipulate the expression until it is identical to the other side.

EXAMPLE 3 Verifying a Trigonometric Identity

Verify the identity $\frac{\sec^2 \theta - 1}{\sec^2 \theta} = \sin^2 \theta$.

SOLUTION

$$\begin{aligned}\frac{\sec^2 \theta - 1}{\sec^2 \theta} &= \frac{\sec^2 \theta}{\sec^2 \theta} - \frac{1}{\sec^2 \theta} && \text{Write as separate fractions.} \\ &= 1 - \left(\frac{1}{\sec \theta}\right)^2 && \text{Simplify.} \\ &= 1 - \cos^2 \theta && \text{Reciprocal identity} \\ &= \sin^2 \theta && \text{Pythagorean identity}\end{aligned}$$

Notice that verifying an identity is not the same as solving an equation. When verifying an identity, you cannot assume that the two sides of the equation are equal because you are trying to verify that they are equal. So, you cannot use any properties of equality, such as adding the same quantity to each side of the equation.

EXAMPLE 4 Verifying a Trigonometric Identity

Verify the identity $\sec x + \tan x = \frac{\cos x}{1 - \sin x}$.

SOLUTION

$$\begin{aligned}\sec x + \tan x &= \frac{1}{\cos x} + \tan x && \text{Reciprocal identity} \\ &= \frac{1}{\cos x} + \frac{\sin x}{\cos x} && \text{Tangent identity} \\ &= \frac{1 + \sin x}{\cos x} && \text{Add fractions.} \\ &= \frac{1 + \sin x}{\cos x} \cdot \frac{1 - \sin x}{1 - \sin x} && \text{Multiply by } \frac{1 - \sin x}{1 - \sin x}. \\ &= \frac{1 - \sin^2 x}{\cos x(1 - \sin x)} && \text{Simplify numerator.} \\ &= \frac{\cos^2 x}{\cos x(1 - \sin x)} && \text{Pythagorean identity} \\ &= \frac{\cos x}{1 - \sin x} && \text{Simplify.}\end{aligned}$$

LOOKING FOR STRUCTURE

To verify the identity, you must introduce $1 - \sin x$ into the denominator. Multiply the numerator and the denominator by $1 - \sin x$ so you get an equivalent expression.



Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Verify the identity.

5. $\cot(-\theta) = -\cot \theta$

7. $\cos x \csc x \tan x = 1$

6. $\csc^2 x(1 - \sin^2 x) = \cot^2 x$

8. $(\tan^2 x + 1)(\cos^2 x - 1) = -\tan^2 x$

9.7 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- WRITING** Describe the difference between a trigonometric identity and a trigonometric equation.
- WRITING** Explain how to use trigonometric identities to determine whether $\sec(-\theta) = \sec \theta$ or $\sec(-\theta) = -\sec \theta$.

Monitoring Progress and Modeling with Mathematics

In Exercises 3–10, find the values of the other five trigonometric functions of θ . (See Example 1.)

- $\sin \theta = \frac{1}{3}, 0 < \theta < \frac{\pi}{2}$
- $\sin \theta = -\frac{7}{10}, \pi < \theta < \frac{3\pi}{2}$
- $\tan \theta = -\frac{3}{7}, \frac{\pi}{2} < \theta < \pi$
- $\cot \theta = -\frac{2}{5}, \frac{\pi}{2} < \theta < \pi$
- $\cos \theta = -\frac{5}{6}, \pi < \theta < \frac{3\pi}{2}$
- $\sec \theta = \frac{9}{4}, \frac{3\pi}{2} < \theta < 2\pi$
- $\cot \theta = -3, \frac{3\pi}{2} < \theta < 2\pi$
- $\csc \theta = -\frac{5}{3}, \pi < \theta < \frac{3\pi}{2}$

In Exercises 11–20, simplify the expression. (See Example 2.)

- $\sin x \cot x$
- $\cos \theta(1 + \tan^2 \theta)$
- $\frac{\sin(-\theta)}{\cos(-\theta)}$
- $\frac{\cos^2 x}{\cot^2 x}$
- $\frac{\cos\left(\frac{\pi}{2} - x\right)}{\csc x}$
- $\sin\left(\frac{\pi}{2} - \theta\right) \sec \theta$
- $\frac{\csc^2 x - \cot^2 x}{\sin(-x) \cot x}$
- $\frac{\cos^2 x \tan^2(-x) - 1}{\cos^2 x}$
- $\frac{\cos\left(\frac{\pi}{2} - \theta\right)}{\csc \theta} + \cos^2 \theta$
- $\frac{\sec x \sin x + \cos\left(\frac{\pi}{2} - x\right)}{1 + \sec x}$

ERROR ANALYSIS In Exercises 21 and 22, describe and correct the error in simplifying the expression.

21.



$$\begin{aligned} 1 - \sin^2 \theta &= 1 - (1 + \cos^2 \theta) \\ &= 1 - 1 - \cos^2 \theta \\ &= -\cos^2 \theta \end{aligned}$$

22.



$$\begin{aligned} \tan x \csc x &= \frac{\cos x}{\sin x} \cdot \frac{1}{\sin x} \\ &= \frac{\cos x}{\sin^2 x} \end{aligned}$$


In Exercises 23–30, verify the identity. (See Examples 3 and 4.)

- $\sin x \csc x = 1$
- $\tan \theta \csc \theta \cos \theta = 1$
- $\cos\left(\frac{\pi}{2} - x\right) \cot x = \cos x$
- $\sin\left(\frac{\pi}{2} - x\right) \tan x = \sin x$
- $\frac{\cos\left(\frac{\pi}{2} - \theta\right) + 1}{1 - \sin(-\theta)} = 1$
- $\frac{\sin^2(-x)}{\tan^2 x} = \cos^2 x$
- $\frac{1 + \cos x}{\sin x} + \frac{\sin x}{1 + \cos x} = 2 \csc x$
- $\frac{\sin x}{1 - \cos(-x)} = \csc x + \cot x$
- USING STRUCTURE** A function f is *odd* when $f(-x) = -f(x)$. A function f is *even* when $f(-x) = f(x)$. Which of the six trigonometric functions are odd? Which are even? Justify your answers using identities and graphs.
- ANALYZING RELATIONSHIPS** As the value of $\cos \theta$ increases, what happens to the value of $\sec \theta$? Explain your reasoning.


33. **MAKING AN ARGUMENT** Your friend simplifies an expression and obtains $\sec x \tan x - \sin x$. You simplify the same expression and obtain $\sin x \tan^2 x$. Are your answers equivalent? Justify your answer.

34. **HOW DO YOU SEE IT?** The figure shows the unit circle and the angle θ .

- Is $\sin \theta$ positive or negative? $\cos \theta$? $\tan \theta$?
- In what quadrant does the terminal side of $-\theta$ lie?
- Is $\sin(-\theta)$ positive or negative? $\cos(-\theta)$? $\tan(-\theta)$?



35. **MODELING WITH MATHEMATICS** A vertical *gnomon* (the part of a sundial that projects a shadow) has height h . The length s of the shadow cast by the gnomon when the angle of the Sun above the horizon is θ can be modeled by the equation below. Show that the equation below is equivalent to $s = h \cot \theta$.



$$s = \frac{h \sin(90^\circ - \theta)}{\sin \theta}$$


36. **THOUGHT PROVOKING** Explain how you can use a trigonometric identity to find all the values of x for which $\sin x = \cos x$.

37. **DRAWING CONCLUSIONS** Static friction is the amount of force necessary to keep a stationary object on a flat surface from moving. Suppose a book weighing W pounds is lying on a ramp inclined at an angle θ . The coefficient of static friction u for the book can be found using the equation $uW \cos \theta = W \sin \theta$.

- Solve the equation for u and simplify the result.
- Use the equation from part (a) to determine what happens to the value of u as the angle θ increases from 0° to 90° .

38. **PROBLEM SOLVING** When light traveling in a medium (such as air) strikes the surface of a second medium (such as water) at an angle θ_1 , the light begins to travel at a different angle θ_2 . This change of direction is defined by Snell's law, $n_1 \sin \theta_1 = n_2 \sin \theta_2$, where n_1 and n_2 are the *indices of refraction* for the two mediums. Snell's law can be derived from the equation

$$\frac{n_1}{\sqrt{\cot^2 \theta_1 + 1}} = \frac{n_2}{\sqrt{\cot^2 \theta_2 + 1}}$$






- Simplify the equation to derive Snell's law.
 - What is the value of n_1 when $\theta_1 = 55^\circ$, $\theta_2 = 35^\circ$, and $n_2 = 2$?
 - If $\theta_1 = \theta_2$, then what must be true about the values of n_1 and n_2 ? Explain when this situation would occur.
39. **WRITING** Explain how transformations of the graph of the parent function $f(x) = \sin x$ support the cofunction identity $\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$.
40. **USING STRUCTURE** Verify each identity.

- $\ln|\sec \theta| = -\ln|\cos \theta|$
- $\ln|\tan \theta| = \ln|\sin \theta| - \ln|\cos \theta|$

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Find the value of x for the right triangle. (Section 9.1)




9.8 Using Sum and Difference Formulas

Essential Question How can you evaluate trigonometric functions of the sum or difference of two angles?

EXPLORATION 1 Deriving a Difference Formula

Work with a partner.

- a. Explain why the two triangles shown are congruent.



- b. Use the Distance Formula to write an expression for d in the first unit circle.
 c. Use the Distance Formula to write an expression for d in the second unit circle.
 d. Write an equation that relates the expressions in parts (b) and (c). Then simplify this equation to obtain a formula for $\cos(a - b)$.

CONSTRUCTING VIABLE ARGUMENTS

To be proficient in math, you need to understand and use stated assumptions, definitions, and previously established results.

EXPLORATION 2 Deriving a Sum Formula

Work with a partner. Use the difference formula you derived in Exploration 1 to write a formula for $\cos(a + b)$ in terms of sine and cosine of a and b . *Hint:* Use the fact that

$$\cos(a + b) = \cos[a - (-b)].$$

EXPLORATION 3 Deriving Difference and Sum Formulas

Work with a partner. Use the formulas you derived in Explorations 1 and 2 to write formulas for $\sin(a - b)$ and $\sin(a + b)$ in terms of sine and cosine of a and b . *Hint:* Use the cofunction identities

$$\sin\left(\frac{\pi}{2} - a\right) = \cos a \text{ and } \cos\left(\frac{\pi}{2} - a\right) = \sin a$$

and the fact that

$$\cos\left[\left(\frac{\pi}{2} - a\right) + b\right] = \sin(a - b) \text{ and } \sin(a + b) = \sin[a - (-b)].$$

Communicate Your Answer

4. How can you evaluate trigonometric functions of the sum or difference of two angles?
5. a. Find the exact values of $\sin 75^\circ$ and $\cos 75^\circ$ using sum formulas. Explain your reasoning.
 b. Find the exact values of $\sin 75^\circ$ and $\cos 75^\circ$ using difference formulas. Compare your answers to those in part (a).

9.8 Lesson

Core Vocabulary

Previous
ratio

What You Will Learn

- ▶ Use sum and difference formulas to evaluate and simplify trigonometric expressions.
- ▶ Use sum and difference formulas to solve trigonometric equations and rewrite real-life formulas.

Using Sum and Difference Formulas

In this lesson, you will study formulas that allow you to evaluate trigonometric functions of the sum or difference of two angles.

Core Concept

Sum and Difference Formulas

Sum Formulas

$$\sin(a + b) = \sin a \cos b + \cos a \sin b$$

$$\cos(a + b) = \cos a \cos b - \sin a \sin b$$

$$\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}$$

Difference Formulas

$$\sin(a - b) = \sin a \cos b - \cos a \sin b$$

$$\cos(a - b) = \cos a \cos b + \sin a \sin b$$

$$\tan(a - b) = \frac{\tan a - \tan b}{1 + \tan a \tan b}$$

In general, $\sin(a + b) \neq \sin a + \sin b$. Similar statements can be made for the other trigonometric functions of sums and differences.

EXAMPLE 1 Evaluating Trigonometric Expressions

Find the exact value of (a) $\sin 15^\circ$ and (b) $\tan \frac{7\pi}{12}$.

SOLUTION

a. $\sin 15^\circ = \sin(60^\circ - 45^\circ)$ Substitute $60^\circ - 45^\circ$ for 15° .

$$= \sin 60^\circ \cos 45^\circ - \cos 60^\circ \sin 45^\circ$$

Difference formula for sine

$$= \frac{\sqrt{3}}{2} \left(\frac{\sqrt{2}}{2} \right) - \frac{1}{2} \left(\frac{\sqrt{2}}{2} \right)$$

Evaluate.

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

Simplify.

▶ The exact value of $\sin 15^\circ$ is $\frac{\sqrt{6} - \sqrt{2}}{4}$. Check this with a calculator.

b. $\tan \frac{7\pi}{12} = \tan \left(\frac{\pi}{3} + \frac{\pi}{4} \right)$ Substitute $\frac{\pi}{3} + \frac{\pi}{4}$ for $\frac{7\pi}{12}$.

$$= \frac{\tan \frac{\pi}{3} + \tan \frac{\pi}{4}}{1 - \tan \frac{\pi}{3} \tan \frac{\pi}{4}}$$

Sum formula for tangent

$$= \frac{\sqrt{3} + 1}{1 - \sqrt{3} \cdot 1}$$

Evaluate.

$$= -2 - \sqrt{3}$$

Simplify.

▶ The exact value of $\tan \frac{7\pi}{12}$ is $-2 - \sqrt{3}$. Check this with a calculator.

Check

$$\begin{aligned} \sin(15^\circ) & \\ & .2588190451 \\ (\sqrt{6} - \sqrt{2})/4 & \\ & .2588190451 \end{aligned}$$

Check

$$\begin{aligned} \tan(7\pi/12) & \\ & -3.732050808 \\ -2 - \sqrt{3} & \\ & -3.732050808 \end{aligned}$$

ANOTHER WAY

You can also use a Pythagorean identity and quadrant signs to find $\sin a$ and $\cos b$.


EXAMPLE 2 Using a Difference Formula

Find $\cos(a - b)$ given that $\cos a = -\frac{4}{5}$ with $\pi < a < \frac{3\pi}{2}$ and $\sin b = \frac{5}{13}$ with $0 < b < \frac{\pi}{2}$.


SOLUTION

Step 1 Find $\sin a$ and $\cos b$.

Because $\cos a = -\frac{4}{5}$ and a is in Quadrant III, $\sin a = -\frac{3}{5}$, as shown in the figure.



Because $\sin b = \frac{5}{13}$ and b is in Quadrant I, $\cos b = \frac{12}{13}$, as shown in the figure.



Step 2 Use the difference formula for cosine to find $\cos(a - b)$.

$$\begin{aligned}\cos(a - b) &= \cos a \cos b + \sin a \sin b && \text{Difference formula for cosine} \\ &= -\frac{4}{5} \left(\frac{12}{13} \right) + \left(-\frac{3}{5} \right) \left(\frac{5}{13} \right) && \text{Evaluate.} \\ &= -\frac{63}{65} && \text{Simplify.}\end{aligned}$$

► The value of $\cos(a - b)$ is $-\frac{63}{65}$.

EXAMPLE 3 Simplifying an Expression

Simplify the expression $\cos(x + \pi)$.

SOLUTION

$$\begin{aligned}\cos(x + \pi) &= \cos x \cos \pi - \sin x \sin \pi && \text{Sum formula for cosine} \\ &= (\cos x)(-1) - (\sin x)(0) && \text{Evaluate.} \\ &= -\cos x && \text{Simplify.}\end{aligned}$$

Monitoring Progress



Help in English and Spanish at BigIdeasMath.com

Find the exact value of the expression.

1. $\sin 105^\circ$
2. $\cos 15^\circ$
3. $\tan \frac{5\pi}{12}$
4. $\cos \frac{\pi}{12}$
5. Find $\sin(a - b)$ given that $\sin a = \frac{8}{17}$ with $0 < a < \frac{\pi}{2}$ and $\cos b = -\frac{24}{25}$ with $\pi < b < \frac{3\pi}{2}$.

Simplify the expression.

6. $\sin(x + \pi)$
7. $\cos(x - 2\pi)$
8. $\tan(x - \pi)$

Solving Equations and Rewriting Formulas

EXAMPLE 4 Solving a Trigonometric Equation

Solve $\sin\left(x + \frac{\pi}{3}\right) + \sin\left(x - \frac{\pi}{3}\right) = 1$ for $0 \leq x < 2\pi$.

SOLUTION

$$\sin\left(x + \frac{\pi}{3}\right) + \sin\left(x - \frac{\pi}{3}\right) = 1$$

Write equation.

$$\sin x \cos \frac{\pi}{3} + \cos x \sin \frac{\pi}{3} + \sin x \cos \frac{\pi}{3} - \cos x \sin \frac{\pi}{3} = 1$$

Use formulas.

$$\frac{1}{2}\sin x + \frac{\sqrt{3}}{2}\cos x + \frac{1}{2}\sin x - \frac{\sqrt{3}}{2}\cos x = 1$$

Evaluate.

$$\sin x = 1$$

Simplify.

► In the interval $0 \leq x < 2\pi$, the solution is $x = \frac{\pi}{2}$.

EXAMPLE 5 Rewriting a Real-Life Formula

The *index of refraction* of a transparent material is the ratio of the speed of light in a vacuum to the speed of light in the material. A triangular prism, like the one shown, can be used to measure the index of refraction using the formula

$$n = \frac{\sin\left(\frac{\theta}{2} + \frac{\alpha}{2}\right)}{\sin \frac{\theta}{2}}.$$

For $\alpha = 60^\circ$, show that the formula can be rewritten as $n = \frac{\sqrt{3}}{2} + \frac{1}{2} \cot \frac{\theta}{2}$.

SOLUTION

$$n = \frac{\sin\left(\frac{\theta}{2} + 30^\circ\right)}{\sin \frac{\theta}{2}}$$

Write formula with $\frac{\alpha}{2} = \frac{60^\circ}{2} = 30^\circ$.

$$= \frac{\sin \frac{\theta}{2} \cos 30^\circ + \cos \frac{\theta}{2} \sin 30^\circ}{\sin \frac{\theta}{2}}$$

Sum formula for sine

$$= \frac{\left(\sin \frac{\theta}{2}\right)\left(\frac{\sqrt{3}}{2}\right) + \left(\cos \frac{\theta}{2}\right)\left(\frac{1}{2}\right)}{\sin \frac{\theta}{2}}$$

Evaluate.

$$= \frac{\frac{\sqrt{3}}{2} \sin \frac{\theta}{2}}{\sin \frac{\theta}{2}} + \frac{\frac{1}{2} \cos \frac{\theta}{2}}{\sin \frac{\theta}{2}}$$


Write as separate fractions.

$$= \frac{\sqrt{3}}{2} + \frac{1}{2} \cot \frac{\theta}{2}$$

Simplify.

ANOTHER WAY

You can also solve the equation by using a graphing calculator. First, graph each side of the original equation. Then use the *intersect* feature to find the x -value(s) where the expressions are equal.



Monitoring Progress  Help in English and Spanish at BigIdeasMath.com

9. Solve $\sin\left(\frac{\pi}{4} - x\right) - \sin\left(x + \frac{\pi}{4}\right) = 1$ for $0 \leq x < 2\pi$.

9.8 Exercises

Dynamic Solutions available at BigIdeasMath.com

Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** Write the expression $\cos 130^\circ \cos 40^\circ - \sin 130^\circ \sin 40^\circ$ as the cosine of an angle.
- WRITING** Explain how to evaluate $\tan 75^\circ$ using either the sum or difference formula for tangent.

Monitoring Progress and Modeling with Mathematics

In Exercises 3–10, find the exact value of the expression.
(See Example 1.)

- $\tan(-15^\circ)$
- $\tan 195^\circ$
- $\sin \frac{23\pi}{12}$
- $\sin(-165^\circ)$
- $\cos 105^\circ$
- $\cos \frac{11\pi}{12}$
- $\tan \frac{17\pi}{12}$
- $\sin\left(-\frac{7\pi}{12}\right)$

In Exercises 11–16, evaluate the expression given that $\cos a = \frac{4}{5}$ with $0 < a < \frac{\pi}{2}$ and $\sin b = -\frac{15}{17}$ with $\frac{3\pi}{2} < b < 2\pi$. (See Example 2.)

- $\sin(a + b)$
- $\sin(a - b)$
- $\cos(a - b)$
- $\cos(a + b)$
- $\tan(a + b)$
- $\tan(a - b)$

In Exercises 17–22, simplify the expression.
(See Example 3.)

- $\tan(x + \pi)$
- $\cos\left(x - \frac{\pi}{2}\right)$
- $\cos(x + 2\pi)$
- $\tan(x - 2\pi)$
- $\sin\left(x - \frac{3\pi}{2}\right)$
- $\tan\left(x + \frac{\pi}{2}\right)$

ERROR ANALYSIS In Exercises 23 and 24, describe and correct the error in simplifying the expression.

23.

$$\begin{aligned} \tan\left(x + \frac{\pi}{4}\right) &= \frac{\tan x + \tan \frac{\pi}{4}}{1 + \tan x \tan \frac{\pi}{4}} \\ &= \frac{\tan x + 1}{1 + \tan x} \\ &= 1 \end{aligned}$$

24.

$$\begin{aligned} \sin\left(x - \frac{\pi}{4}\right) &= \sin \frac{\pi}{4} \cos x - \cos \frac{\pi}{4} \sin x \\ &= \frac{\sqrt{2}}{2} \cos x - \frac{\sqrt{2}}{2} \sin x \\ &= \frac{\sqrt{2}}{2} (\cos x - \sin x) \end{aligned}$$

25. What are the solutions of the equation $2 \sin x - 1 = 0$ for $0 \leq x < 2\pi$?

- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{6}$
(C) $\frac{2\pi}{3}$ (D) $\frac{5\pi}{6}$

26. What are the solutions of the equation $\tan x + 1 = 0$ for $0 \leq x < 2\pi$?

- (A) $\frac{\pi}{4}$ (B) $\frac{3\pi}{4}$
(C) $\frac{5\pi}{4}$ (D) $\frac{7\pi}{4}$

In Exercises 27–32, solve the equation for $0 \leq x < 2\pi$.
(See Example 4.)

27. $\sin\left(x + \frac{\pi}{2}\right) = \frac{1}{2}$ 28. $\tan\left(x - \frac{\pi}{4}\right) = 0$

29. $\cos\left(x + \frac{\pi}{6}\right) - \cos\left(x - \frac{\pi}{6}\right) = 1$

30. $\sin\left(x + \frac{\pi}{4}\right) + \sin\left(x - \frac{\pi}{4}\right) = 0$

31. $\tan(x + \pi) - \tan(\pi - x) = 0$


32. $\sin(x + \pi) + \cos(x + \pi) = 0$

33. **USING EQUATIONS** Derive the cofunction identity $\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$ using the difference formula for sine.

34. **MAKING AN ARGUMENT** Your friend claims it is possible to use the difference formula for tangent to derive the cofunction identity $\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta$. Is your friend correct? Explain your reasoning.

35. **MODELING WITH MATHEMATICS** A photographer is at a height h taking aerial photographs with a 35-millimeter camera. The ratio of the image length WQ to the length NA of the actual object is given by the formula

$$\frac{WQ}{NA} = \frac{35 \tan(\theta - t) + 35 \tan t}{h \tan \theta}$$



where θ is the angle between the vertical line perpendicular to the ground and the line from the camera to point A and t is the tilt angle of the film. When $t = 45^\circ$, show that the formula can be rewritten as $\frac{WQ}{NA} = \frac{70}{h(1 + \tan \theta)}$. (See Example 5.)

36. **MODELING WITH MATHEMATICS** When a wave travels through a taut string, the displacement y of each point on the string depends on the time t and the point's position x . The equation of a *standing wave* can be obtained by adding the displacements of two waves traveling in opposite directions. Suppose a standing wave can be modeled by the formula
- $$y = A \cos\left(\frac{2\pi t}{3} - \frac{2\pi x}{5}\right) + A \cos\left(\frac{2\pi t}{3} + \frac{2\pi x}{5}\right).$$
- When $t = 1$, show that the formula can be rewritten as
- $$y = -A \cos \frac{2\pi x}{5}.$$


37. **MODELING WITH MATHEMATICS** The busy signal on a touch-tone phone is a combination of two tones with frequencies of 480 hertz and 620 hertz. The individual tones can be modeled by the equations:

480 hertz: $y_1 = \cos 960\pi t$


620 hertz: $y_2 = \cos 1240\pi t$

The sound of the busy signal can be modeled by $y_1 + y_2$. Show that $y_1 + y_2 = 2 \cos 1100\pi t \cos 140\pi t$.

38. **HOW DO YOU SEE IT?** Explain how to use the figure to solve the equation $\sin\left(x + \frac{\pi}{4}\right) - \sin\left(\frac{\pi}{4} - x\right) = 0$ for $0 \leq x < 2\pi$.



39. **MATHEMATICAL CONNECTIONS** The figure shows the acute angle of intersection, $\theta_2 - \theta_1$, of two lines with slopes m_1 and m_2 .



- Use the difference formula for tangent to write an equation for $\tan(\theta_2 - \theta_1)$ in terms of m_1 and m_2 .
- Use the equation from part (a) to find the acute angle of intersection of the lines $y = x - 1$ and $y = \left(\frac{1}{\sqrt{3} - 2}\right)x + \frac{4 - \sqrt{3}}{2 - \sqrt{3}}$.

40. **THOUGHT PROVOKING** Rewrite each function. Justify your answers.

- Write $\sin 3x$ as a function of $\sin x$.
- Write $\cos 3x$ as a function of $\cos x$.
- Write $\tan 3x$ as a function of $\tan x$.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Solve the equation. Check your solution(s). (Section 7.5)

41. $1 - \frac{9}{x-2} = -\frac{7}{2}$

42. $\frac{12}{x} + \frac{3}{4} = \frac{8}{x}$

43. $\frac{2x-3}{x+1} = \frac{10}{x^2-1} + 5$

9.5–9.8 What Did You Learn?

Core Vocabulary

frequency, *p.* 506

sinusoid, *p.* 507

trigonometric identity, *p.* 514

Core Concepts

Section 9.5

Characteristics of $y = \tan x$ and $y = \cot x$, *p.* 498

Period and Vertical Asymptotes of $y = a \tan bx$ and $y = a \cot bx$, *p.* 499

Characteristics of $y = \sec x$ and $y = \csc x$, *p.* 500

Section 9.6

Frequency, *p.* 506

Writing Trigonometric Functions, *p.* 507

Using Technology to Find Trigonometric Models, *p.* 509

Section 9.7

Fundamental Trigonometric Identities, *p.* 514

Section 9.8

Sum and Difference Formulas, *p.* 520

Trigonometric Equations and Real-Life Formulas, *p.* 522

Mathematical Practices

1. Explain why the relationship between θ and d makes sense in the context of the situation in Exercise 43 on page 503.
2. How can you use definitions to relate the slope of a line with the tangent of an angle in Exercise 39 on page 524?

Performance Task

Lightening the Load

You need to move a heavy table across the room. What is the easiest way to move it? Should you push it? Should you tie a rope around one leg of the table and pull it? How can trigonometry help you make the right decision?

To explore the answers to these questions and more, go to BigIdeasMath.com.



9 Chapter Review


Dynamic Solutions available at BigIdeasMath.com

9.1 Right Triangle Trigonometry (pp. 461–468)

Evaluate the six trigonometric functions of the angle θ .

From the Pythagorean Theorem, the length of the hypotenuse is


$$\begin{aligned}\text{hyp.} &= \sqrt{6^2 + 8^2} \\ &= \sqrt{100} \\ &= 10.\end{aligned}$$



Using $\text{adj.} = 8$, $\text{opp.} = 6$, and $\text{hyp.} = 10$, the values of the six trigonometric functions of θ are:

$$\begin{aligned}\sin \theta &= \frac{\text{opp.}}{\text{hyp.}} = \frac{6}{10} = \frac{3}{5} & \cos \theta &= \frac{\text{adj.}}{\text{hyp.}} = \frac{8}{10} = \frac{4}{5} & \tan \theta &= \frac{\text{opp.}}{\text{adj.}} = \frac{6}{8} = \frac{3}{4} \\ \csc \theta &= \frac{\text{hyp.}}{\text{opp.}} = \frac{10}{6} = \frac{5}{3} & \sec \theta &= \frac{\text{hyp.}}{\text{adj.}} = \frac{10}{8} = \frac{5}{4} & \cot \theta &= \frac{\text{adj.}}{\text{opp.}} = \frac{8}{6} = \frac{4}{3}\end{aligned}$$

1. In a right triangle, θ is an acute angle and $\cos \theta = \frac{6}{11}$. Evaluate the other five trigonometric functions of θ .
2. The shadow of a tree measures 25 feet from its base. The angle of elevation to the Sun is 31° . How tall is the tree?



9.2 Angles and Radian Measure (pp. 469–476)

Convert the degree measure to radians or the radian measure to degrees.

a. 110°

$$\begin{aligned}110^\circ &= 110 \cancel{\text{degrees}} \left(\frac{\pi \text{ radians}}{180 \cancel{\text{degrees}}} \right) \\ &= \frac{11\pi}{18}\end{aligned}$$

b. $\frac{7\pi}{12}$

$$\begin{aligned}\frac{7\pi}{12} &= \frac{7\pi}{12} \cancel{\text{radians}} \left(\frac{180^\circ}{\pi \cancel{\text{radians}}} \right) \\ &= 105^\circ\end{aligned}$$

3. Find one positive angle and one negative angle that are coterminal with 382° .

Convert the degree measure to radians or the radian measure to degrees.

4. 30°

5. 225°

6. $\frac{3\pi}{4}$

7. $\frac{5\pi}{3}$

8. A sprinkler system on a farm rotates 140° and sprays water up to 35 meters. Draw a diagram that shows the region that can be irrigated with the sprinkler. Then find the area of the region.


9.3 Trigonometric Functions of Any Angle (pp. 477–484)

Evaluate $\csc 210^\circ$.


The reference angle is $\theta' = 210^\circ - 180^\circ = 30^\circ$. The cosecant function is negative in Quadrant III, so $\csc 210^\circ = -\csc 30^\circ = -2$.

Evaluate the six trigonometric functions of θ .


9.



10.



11.



Evaluate the function without using a calculator.

12. $\tan 330^\circ$

13. $\sec(-405^\circ)$

14. $\sin \frac{13\pi}{6}$

15. $\sec \frac{11\pi}{3}$

9.4 Graphing Sine and Cosine Functions (pp. 485–494)


Identify the amplitude and period of $g(x) = \frac{1}{2} \sin 2x$. Then graph the function and describe the graph of g as a transformation of the graph of $f(x) = \sin x$.

The function is of the form $g(x) = a \sin bx$, where $a = \frac{1}{2}$ and $b = 2$. So, the amplitude is $a = \frac{1}{2}$ and the period is $\frac{2\pi}{b} = \frac{2\pi}{2} = \pi$.

Intercepts: $(0, 0); \left(\frac{1}{2} \cdot \pi, 0\right) = \left(\frac{\pi}{2}, 0\right); (\pi, 0)$

Maximum: $\left(\frac{1}{4} \cdot \pi, \frac{1}{2}\right) = \left(\frac{\pi}{4}, \frac{1}{2}\right)$

Minimum: $\left(\frac{3}{4} \cdot \pi, -\frac{1}{2}\right) = \left(\frac{3\pi}{4}, -\frac{1}{2}\right)$



► The graph of g is a vertical shrink by a factor of $\frac{1}{2}$ and a horizontal shrink by a factor of $\frac{1}{2}$ of the graph of f .

Identify the amplitude and period of the function. Then graph the function and describe the graph of g as a transformation of the graph of the parent function.

16. $g(x) = 8 \cos x$

17. $g(x) = 6 \sin \pi x$

18. $g(x) = \frac{1}{4} \cos 4x$

Graph the function.

19. $g(x) = \cos(x + \pi) + 2$

20. $g(x) = -\sin x - 4$

21. $g(x) = 2 \sin\left(x + \frac{\pi}{2}\right)$

9.5 Graphing Other Trigonometric Functions (pp. 497–504)


- a. Graph one period of $g(x) = 7 \cot \pi x$. Describe the graph of g as a transformation of the graph of $f(x) = \cot x$.

The function is of the form $g(x) = a \cot bx$, where $a = 7$ and $b = \pi$. So, the period is $\frac{\pi}{|b|} = \frac{\pi}{\pi} = 1$.

$$\text{Intercepts: } \left(\frac{\pi}{2b}, 0\right) = \left(\frac{\pi}{2\pi}, 0\right) = \left(\frac{1}{2}, 0\right)$$

$$\text{Asymptotes: } x = 0; x = \frac{\pi}{|b|} = \frac{\pi}{\pi}, \text{ or } x = 1$$

$$\begin{aligned} \text{Halfway points: } \left(\frac{\pi}{4b}, a\right) &= \left(\frac{\pi}{4\pi}, 7\right) = \left(\frac{1}{4}, 7\right); \\ \left(\frac{3\pi}{4b}, -a\right) &= \left(\frac{3\pi}{4\pi}, -7\right) = \left(\frac{3}{4}, -7\right) \end{aligned}$$



- The graph of g is a vertical stretch by a factor of 7 and a horizontal shrink by a factor of $\frac{1}{\pi}$ of the graph of f .


- b. Graph one period of $g(x) = 9 \sec x$. Describe the graph of g as a transformation of the graph of $f(x) = \sec x$.

Step 1 Graph the function $y = 9 \cos x$.

The period is $\frac{2\pi}{1} = 2\pi$.

Step 2 Graph asymptotes of g . Because the asymptotes of g occur when $9 \cos x = 0$, graph $x = -\frac{\pi}{2}$, $x = \frac{\pi}{2}$, and $x = \frac{3\pi}{2}$.

Step 3 Plot the points on g , such as $(0, 9)$ and $(\pi, -9)$. Then use the asymptotes to sketch the curve.



- The graph of g is a vertical stretch by a factor of 9 of the graph of f .

Graph one period of the function. Describe the graph of g as a transformation of the graph of its parent function.

22. $g(x) = \tan \frac{1}{2}x$

23. $g(x) = 2 \cot x$

24. $g(x) = 4 \tan 3\pi x$

Graph the function.

25. $g(x) = 5 \csc x$

26. $g(x) = \sec \frac{1}{2}x$

27. $g(x) = 5 \sec \pi x$

28. $g(x) = \frac{1}{2} \csc \frac{\pi}{4}x$

9.6 Modeling with Trigonometric Functions (pp. 505–512)

Write a function for the sinusoid shown.

Step 1 Find the maximum and minimum values. From the graph, the maximum value is 3 and the minimum value is -1 .

Step 2 Identify the vertical shift, k . The value of k is the mean of the maximum and minimum values.

$$k = \frac{(\text{maximum value}) + (\text{minimum value})}{2} = \frac{3 + (-1)}{2} = \frac{2}{2} = 1$$

Step 3 Decide whether the graph should be modeled by a sine or cosine function. Because the graph crosses the midline $y = 1$ on the y -axis and then decreases to its minimum value, the graph is a sine curve with a reflection in the x -axis and no horizontal shift. So, $h = 0$.

Step 4 Find the amplitude and period.


The period is $\frac{2\pi}{3} = \frac{2\pi}{b}$. So, $b = 3$.

The amplitude is

$$|a| = \frac{(\text{maximum value}) - (\text{minimum value})}{2} = \frac{3 - (-1)}{2} = \frac{4}{2} = 2.$$


Because the graph is a reflection in the x -axis, $a < 0$. So, $a = -2$.

► The function is $y = -2 \sin 3x + 1$.




Write a function for the sinusoid.

29.



30.



31. You put a reflector on a spoke of your bicycle wheel. The highest point of the reflector is 25 inches above the ground, and the lowest point is 2 inches. The reflector makes 1 revolution per second. Write a model for the height h (in inches) of a reflector as a function of time t (in seconds) given that the reflector is at its lowest point when $t = 0$.

32. The table shows the monthly precipitation P (in inches) for Bismarck, North Dakota, where $t = 1$ represents January. Write a model that gives P as a function of t and interpret the period of its graph.

t	1	2	3	4	5	6	7	8	9	10	11	12
P	0.5	0.5	0.9	1.5	2.2	2.6	2.6	2.2	1.6	1.3	0.7	0.4

9.7 Using Trigonometric Identities (pp. 513–518)

Verify the identity $\frac{\cot^2 \theta}{\csc \theta} = \csc \theta - \sin \theta$.

$$\begin{aligned}\frac{\cot^2 \theta}{\csc \theta} &= \frac{\csc^2 \theta - 1}{\csc \theta} \\ &= \frac{\csc^2 \theta}{\csc \theta} - \frac{1}{\csc \theta} \\ &= \csc \theta - \frac{1}{\csc \theta} \\ &= \csc \theta - \sin \theta\end{aligned}$$

Pythagorean identity

Write as separate fractions.

Simplify.

Reciprocal identity

Simplify the expression.

33. $\cot^2 x - \cot^2 x \cos^2 x$

34. $\frac{(\sec x + 1)(\sec x - 1)}{\tan x}$

35. $\sin\left(\frac{\pi}{2} - x\right) \tan x$

Verify the identity.

36. $\frac{\cos x \sec x}{1 + \tan^2 x} = \cos^2 x$

37. $\tan\left(\frac{\pi}{2} - x\right) \cot x = \csc^2 x - 1$

9.8 Using Sum and Difference Formulas (pp. 519–524)

Find the exact value of $\sin 105^\circ$.

$$\begin{aligned}\sin 105^\circ &= \sin(45^\circ + 60^\circ) \\ &= \sin 45^\circ \cos 60^\circ + \cos 45^\circ \sin 60^\circ \\ &= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} + \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} \\ &= \frac{\sqrt{2} + \sqrt{6}}{4}\end{aligned}$$

Substitute $45^\circ + 60^\circ$ for 105° .

Sum formula for sine

Evaluate.

Simplify.

► The exact value of $\sin 105^\circ$ is $\frac{\sqrt{2} + \sqrt{6}}{4}$.

Find the exact value of the expression.

38. $\sin 75^\circ$

39. $\tan(-15^\circ)$

40. $\cos \frac{\pi}{12}$

41. Find $\tan(a + b)$, given that $\tan a = \frac{1}{4}$ with $\pi < a < \frac{3\pi}{2}$ and $\tan b = \frac{3}{7}$ with $0 < b < \frac{\pi}{2}$.

Solve the equation for $0 \leq x < 2\pi$.

42. $\cos\left(x + \frac{3\pi}{4}\right) + \cos\left(x - \frac{3\pi}{4}\right) = 1$



43. $\tan(x + \pi) + \cos\left(x + \frac{\pi}{2}\right) = 0$

9 Chapter Test

Verify the identity.

- $\frac{\cos^2 x + \sin^2 x}{1 + \tan^2 x} = \cos^2 x$
- $\frac{1 + \sin x}{\cos x} + \frac{\cos x}{1 + \sin x} = 2 \sec x$
- $\cos\left(x + \frac{3\pi}{2}\right) = \sin x$
- Evaluate $\sec(-300^\circ)$ without using a calculator.

Write a function for the sinusoid.





Graph the function. Then describe the graph of g as a transformation of the graph of its parent function.

- $g(x) = -4 \tan 2x$
- $g(x) = -2 \cos \frac{1}{3}x + 3$
- $g(x) = 3 \csc \pi x$

Convert the degree measure to radians or the radian measure to degrees. Then find one positive angle and one negative angle that are coterminal with the given angle.


- -50°
- $\frac{4\pi}{5}$
- $\frac{8\pi}{3}$
- Find the arc length and area of a sector with radius $r = 13$ inches and central angle $\theta = 40^\circ$.

Evaluate the six trigonometric functions of the angle θ .



16. In which quadrant does the terminal side of θ lie when $\cos \theta < 0$ and $\tan \theta > 0$? Explain.

17. How tall is the building? Justify your answer.



18. The table shows the average daily high temperatures T (in degrees Fahrenheit) in Baltimore, Maryland, where $m = 1$ represents January. Write a model that gives T as a function of m and interpret the period of its graph.

m	1	2	3	4	5	6	7	8	9	10	11	12
T	41	45	54	65	74	83	87	85	78	67	56	45

9 Cumulative Assessment

1. Which expressions are equivalent to 1?

$$\tan x \sec x \cos x$$

$$\sin^2 x + \cos^2 x$$

$$\frac{\cos^2(-x) \tan^2 x}{\sin^2(-x)}$$

$$\cos\left(\frac{\pi}{2} - x\right) \csc x$$


2. Which rational expression represents the ratio of the perimeter to the area of the playground shown in the diagram?

(A) $\frac{9}{7x}$

(B) $\frac{11}{14x}$

(C) $\frac{1}{x}$

(D) $\frac{1}{2x}$



3. The chart shows the average monthly temperatures (in degrees Fahrenheit) and the gas usages (in cubic feet) of a household for 12 months.

- a. Use a graphing calculator to find trigonometric models for the average temperature y_1 as a function of time and the gas usage y_2 (in thousands of cubic feet) as a function of time. Let $t = 1$ represent January.

- b. Graph the two regression equations in the same coordinate plane on your graphing calculator. Describe the relationship between the graphs.

January	February	March	April
32°F	21°F	15°F	22°F
20,000 ft ³	27,000 ft ³	23,000 ft ³	22,000 ft ³
May	June	July	August
35°F	49°F	62°F	78°F
21,000 ft ³	14,000 ft ³	8,000 ft ³	9,000 ft ³
September	October	November	December
71°F	63°F	55°F	40°F
13,000 ft ³	15,000 ft ³	19,000 ft ³	23,000 ft ³

4. Evaluate each logarithm using $\log_2 5 \approx 2.322$ and $\log_2 3 \approx 1.585$, if necessary. Then order the logarithms by value from least to greatest.

a. $\log 1000$

b. $\log_2 15$

c. $\ln e$


d. $\log_2 9$

e. $\log_2 \frac{5}{3}$

f. $\log_2 1$


5. Which function is *not* represented by the graph?

- (A) $y = 5 \sin x$
 (B) $y = 5 \cos\left(\frac{\pi}{2} - x\right)$
 (C) $y = 5 \cos\left(x + \frac{\pi}{2}\right)$
 (D) $y = -5 \sin(x + \pi)$




6. Complete each statement with $<$ or $>$ so that each statement is true.

- a. θ 3 radians
 b. $\tan \theta$ 0
 c. θ' 45°



7. Use the Rational Root Theorem and the graph to find all the real zeros of the function $f(x) = 2x^3 - x^2 - 13x - 6$.



8. Your friend claims -210° is coterminal with the angle $\frac{5\pi}{6}$. Is your friend correct? Explain your reasoning.
9. Company A and Company B offer the same starting annual salary of \$20,000. Company A gives a \$1000 raise each year. Company B gives a 4% raise each year.
- Write rules giving the salaries a_n and b_n for your n th year of employment at Company A and Company B, respectively. Tell whether the sequence represented by each rule is *arithmetic*, *geometric*, or *neither*.
 - Graph each sequence in the same coordinate plane.
 - Under what conditions would you choose to work for Company B?
 - After 20 years of employment, compare your total earnings.